

Q1W5-Ph.-H.W-Work and Energy

Multiple Choice

Identify the choice that best completes the statement or answers the question.

- ____ 1. In which of the following sentences is *work* used in the scientific sense of the word?
 - a. Holding a heavy box requires a lot of work.
 - b. A scientist works on an experiment in the laboratory.
 - c. Sam and Rachel pushed hard, but they could do no work on the car.
 - d. John learned that shoveling snow is hard work.
- ____ 2. In which of the following sentences is *work* used in the everyday sense of the word?
 - a. Lifting a heavy bucket involves doing work on the bucket.
 - b. The force of friction usually does negative work.
 - c. Sam and Rachel worked hard pushing the car.
 - d. Work is a physical quantity.
- ____ 3. A force does work on an object if a component of the force
 - a. is perpendicular to the displacement of the object.
 - b. is parallel to the displacement of the object.
 - c. perpendicular to the displacement of the object moves the object along a path that returns the object to its starting position.
 - d. parallel to the displacement of the object moves the object along a path that returns the object to its starting position.
- ____ 4. Work is done when
 - a. the displacement is not zero.
 - b. the displacement is zero.
 - c. the force is zero.
 - d. the force and displacement are perpendicular.
- ____ 5. What is the common formula for work? Assume that W is the work, F is a constant force, Δv is the change in velocity, and d is the displacement.

| | |
|--------------------|-------------------------|
| a. $W = F\Delta v$ | c. $W = F\frac{d^2}{2}$ |
| b. $W = Fd$ | d. $W = F \cdot d$ |
- ____ 6. In which of the following scenarios is work done?
 - a. A weightlifter holds a barbell overhead for 2.5 s.
 - b. A construction worker carries a heavy beam while walking at constant speed along a flat surface.
 - c. A car decelerates while traveling on a flat stretch of road.
 - d. A student holds a spring in a compressed position.
- ____ 7. In which of the following scenarios is no net work done?
 - a. A car accelerates down a hill.
 - b. A car travels at constant speed on a flat road.
 - c. A car decelerates on a flat road.
 - d. A car decelerates as it travels up a hill.
- ____ 8. A child moving at constant velocity carries a 2 N ice-cream cone 1 m across a level surface. What is the net work done on the ice-cream cone?

| | |
|----------|---------|
| a. 0 J | c. 2 J |
| b. 0.5 J | d. 20 J |
- ____ 9. A worker does 25 J of work lifting a bucket, then sets the bucket back down in the same place. What is the total net work done on the bucket?

| | |
|----------|---------|
| a. -25 J | c. 25 J |
| b. 0 J | d. 50 J |

Problem

46. How much work is done on a bookshelf being pulled 4.00 m at an angle of 35.0° from the horizontal? The magnitude of the component of the force that does the work is 87.0 N.

A- 338 J
B- 348 J
C- 358 J
D- 368 J

47. A worker pushes a box with a horizontal force of 40.0 N over a level distance of 4.0 m. If a frictional force of 27 N acts on the box in a direction opposite to that of the worker, what network is done on the box?

A- 46 J
B- 48 J
C- 50 J
D- 52 J

48. A flight attendant pulls a 60.0 N flight bag a distance of 239.0 m along a level airport floor at a constant speed. A 21.0 N force is exerted on the bag at an angle of 66.0° above the horizontal. How much work is done on the flight bag?

A- 2040 J
B- 2540 J
C- 3040 J
D- 3540 J

49. A hill is 132 m long and makes an angle of 12.0° with the horizontal. As a 54 kg jogger runs up the hill, how much work does the jogger do against gravity?

A- 1.1×10^4 J
B- 1.2×10^4 J
C- 1.3×10^4 J
D- 1.5×10^4 J

50. A professional skier starts from rest and reaches a speed of 48 m/s on a ski slope angled 22.0° above the horizontal. Using the work-kinetic energy theorem and disregarding friction, find the minimum distance along the slope the skier would have to travel in order to reach this speed.

A- 310 m
B- 320 m
C- 330 m
D- 340 m

51. A 31.0 kg crate, initially at rest, slides down a ramp 2.6 m long and inclined at an angle of 14.0° with the horizontal. Using the work-kinetic energy theorem and disregarding friction, find the velocity of the crate at the bottom of the ramp. ($g = 9.81 \text{ m/s}^2$)

- A- 3.5 m/s
- B- 3.7 m/s
- C- 3.8 m/s
- D- 3.9 m/s

52. A child riding a bicycle has a total mass of 49.0 kg. The child approaches the top of a hill that is 15.0 m high and 106.0 m long at 14.0 m/s. If the force of friction between the bicycle and the hill is 22.0 N, what is the child's velocity at the bottom of the hill? (Assume no air resistance and that $g = 9.81 \text{ m/s}^2$.)

- A- 18.9 m/s
- B- 19.9 m/s
- C- 21.9 m/s
- D- 23.9 m/s

53. A skier with a mass of 84 kg hits a ramp of snow at 32 m/s and becomes airborne. At the highest point of flight, the skier is 4.7 m above the ground. What is the skier's gravitational potential energy at this point?

- A- $3.3 \times 10^3 \text{ J}$
- B- $3.5 \times 10^3 \text{ J}$
- C- $3.7 \times 10^3 \text{ J}$
- D- $3.9 \times 10^3 \text{ J}$

54. A 2.74 g coin, which has zero potential energy at the surface, is dropped into a 12.2 m well. After the coin comes to a stop in the mud, what is its potential energy with respect to the surface?

- A- -0.208 J
- B- -0.258 J
- C- -0.328 J
- D- -0.358 J

55. A $3.62 \times 10^2 \text{ N}$ crate is pushed to the top of a 2.53 m ramp, which is inclined at 15.0° with the horizontal. What is the potential energy of the crate?

- A- 277 J
- B- 257 J
- C- 237 J
- D- 217 J

56. A 53.0 N crate is pulled up a 5.6 m inclined plane at a constant velocity. If the plane is inclined at an angle of 34.0° to the horizontal and there is a constant force of friction of 15.0 N between the crate and the surface, what is the net gain in potential energy by the crate?

- A- 160 J
- B- 150 J
- C- 140 J
- D- 130 J

57. A 37 kg child on roller skates, initially at rest, rolls 2.0 m down an incline at an angle of 17.0° with the horizontal. If there is no friction between incline and skates, what is the kinetic energy of the child at the bottom of the incline? ($g = 9.81 \text{ m/s}^2$)

- A- 210 J
- B- 205 J
- C- 200 J
- D- 190 J

58. On a given occasion, Old Faithful geyser in Yellowstone National Park shoots water to a height of 47.1 m. With what velocity does the water leave the ground during this eruption? (Assume no air resistance and that $g = 9.81 \text{ m/s}^2$.)

- A- 25.4 m/s
- B- 30.4 m/s
- C- 35.4 m/s
- D- 40.4 m/s

59. A pole vaulter clears 6.41 m. With what velocity does the vaulter strike the mat in the landing area? (Assume no air resistance and that $g = 9.81 \text{ m/s}^2$.)

- A- 9.2 m/s
- B- 10.2 m/s
- C- 11.2 m/s
- D- 12.2 m/s

60. A bobsled zips down an ice track, starting from rest at the top of a hill with a vertical height of 170 m. Disregarding friction, what is the velocity of the bobsled at the bottom of the hill? ($g = 9.81 \text{ m/s}^2$)

- A- 55 m/s
- B- 57 m/s
- C- 59 m/s
- D- 61 m/s

61. A pendulum with a mass of 4.0 kg is released from a height of 2.9 cm above the height of its resting position. How fast will the pendulum be moving when it passes through the lowest point of its swing?

- A- 0.25 m/s
- B- 0.35 m/s
- C- 0.55 m/s
- D- 0.75 m/s

62. What is the average power output of a weightlifter who can lift 260 kg to a height of 1.9 m in 1.8 s?

- A- 2.1 kW
- B- 2.3 kW
- C- 2.5 kW
- D- 2.7 kW

63. A jet engine develops 1.1×10^5 N of thrust to move an airplane forward at a speed of 9.3×10^2 km/h. What is the power output of the engine?

- A- 21 MW
- B- 23 MW
- C- 25 MW
- D- 29 MW

64. A 1.71×10^3 kg sports car accelerates from rest to 25.8 m/s in 7.41 s. What is the average power output of the automobile engine?

- A- 76.8 kW
- B- 74.8 kW
- C- 72.8 kW
- D- 70.8 kW

65. Water flows over a section of Niagara Falls at a rate of 1.30×10^6 kg/s and falls 49.5 m. What is the power of the waterfall?

- A- 621 MW
- B- 631 MW
- C- 641 MW
- D- 651 MW

=====