Trigonometric Functions

_			
2 I.			
in C	;nap	ter 3,	1
stu	died	expon	e

and logarithmic

transcendental

functions.

functions, which are two types of

ential

Then

CHAPTER

Now

- 🧕 In Chapter 4, you will:
 - Use trigonometric functions to solve right triangles.
 - Find values of trigonometric functions for any angle.
 - Graph trigonometric and inverse trigonometric functions.

SATELLITE NAVIGATION Satellite navigation systems operate by receiving signals from satellites in orbit, determining the distance to each of the satellites, and then using trigonometry to establish the location on Earth's surface. These techniques are also used when navigating cars, planes, ships, and spacecraft.

.

PREREAD Use the prereading strategy of previewing to make two or three predictions of what Chapter 4 is about.



Why?

Get Ready for the Chapter

Diagnose Readiness You have two options for checking Prerequisite Skills.



QuickCheck

Find the missing value in each figure. (Prerequisite Skill)



Determine whether each of the following could represent the measures of three sides of a triangle. Write *yes* or *no*. (Prerequisite Skill)

5. 4, 8, 12 **6.** 12, 15, 18

7. ALGEBRA The sides of a triangle have lengths x, x + 17, and 25. If the length of the longest side is 25, what value of x makes the triangle a right triangle? (Prerequisite Skill)

Find the equations of any vertical or horizontal asymptotes. (Lesson 2-5)

8.
$$f(x) = \frac{x^2 - 4}{x^2 + 8}$$
 9. $h(x) = \frac{x^3 - 27}{x + 5}$

10.
$$f(x) = \frac{x(x-1)^2}{(x-2)(x+4)}$$
 11. $g(x) = \frac{x+5}{(x-3)(x-5)}$

12.
$$h(x) = \frac{x^2 + x - 20}{x + 5}$$
 13. $f(x) = \frac{2x^2 + 5x - 12}{2x - 3}$



ewVocabulary		the se
English		Español
trignometric functions	p. 220	funciones trigonométricas
sine	p. 220	seno
cosine	p. 220	coseno
tangent	p. 220	tangente
cosecant	p. 220	cosecant
secant	p. 220	secant
cotangent	p. 220	función recíproca
reciprocal function	p. 220	cotangente
inverse sine	p. 223	seno inverso
inverse cosine	p. 223	coseno inverso
inverse tangent	p. 223	tangente inversa
radian	p. 232	radian
coterminal angles	p. 234	ángulos coterminales
reference angle	p. 244	ángulo de referencia
unit circle	p. 247	círculo de unidad
circular function	p. 248	función circular
period	p. 250	período
sinusoid	p. 256	sinusoid
amplitude	p. 257	amplitud
frequency	p. 260	frecuencia
phase shift	p. 261	cambio de fase
Law of Sines	p. 291	ley de senos
Law of Cosines	p. 295	lev de cosenos

ReviewVocabulary

N



reflection p. 48 reflexion the mirror image of the graph of a function with respect to a specific line

dilation p. 49 homotecia a nonrigid transformation that has the effect of compressing (shrinking) or expanding (enlarging) the graph of a function vertically or horizontally



Vertical dilation

Right Triangle Trigonometry Whv? Now Then You evaluated Find values of Large helium-filled balloons are a tradition of many holiday parades. functions. trigonometric Long cables attached to the balloon are used by volunteers to lead functions for acute the balloon along the parade route. (Lesson 1-1) angles of right Suppose two of these cables are attached to a balloon at the same triangles. point, and the volunteers holding these cables stand so that the ends Solve right triangles. of the cables lie in the same vertical plane. If you know the measure of the angle that each cable makes with the ground and the distance between the volunteers, you can use right triangle trigonometry to find the height of the balloon above the ground. abc Values of Trigonometric Ratios The word trigonometry means triangle measure. In this **NewVocabulary** trigonometric ratios chapter, you will study trigonometry as the relationships among the sides and angles of triangles and as a set of functions defined on the real number system. In this lesson, you will study trigonometric functions right triangle trigonometry. sine cosine Using the side measures of a right triangle and a reference angle labeled θ (the Greek letter theta), tangent we can form the six **trigonometric ratios** that define six **trigonometric functions**. cosecant secant cotangent reciprocal function KeyConcept Trigonometric Functions inverse trigonometric function Let θ be an acute angle in a right triangle and the abbreviations

opp, adj, and hyp refer to the length of the side opposite θ , the

respectively.

sine $(\theta) = \sin \theta = \frac{\operatorname{opp}}{\operatorname{hyp}}$

 $\frac{\text{cosine}}{(\theta)} = \cos \theta = \frac{\text{adj}}{(\theta)}$

tangent $(\theta) = \tan \theta = \frac{\operatorname{opp}}{\operatorname{adi}}$

length of the side adjacent to θ , and the length of the hypotenuse,

Then the six trigonometric functions of θ are defined as follows.

function inverse sine inverse cosine inverse tangent angle of elevation angle of depression solve a right triangle

> The cosecant, secant, and cotangent functions are called **reciprocal functions** because their ratios are reciprocals of the sine, cosine, and tangent ratios, respectively. Therefore, the following statements are true. $\csc \theta = \frac{1}{\sin \theta} \qquad \sec \theta = \frac{1}{\cos \theta} \qquad \cot \theta = \frac{1}{\tan \theta}$

From the definitions of the sine, cosine, tangent, and cotangent functions, you can also derive the following relationships. You will prove these relationships in Exercise 83.

$$\tan \theta = \frac{\sin \theta}{\cos \theta}$$
 and $\cot \theta = \frac{\cos \theta}{\sin \theta}$

hvp

adj

opp

 $\frac{\text{cosecant}}{\text{cosecant}}(\theta) = \csc \theta = \frac{\text{hyp}}{\text{opp}}$

 $\frac{\text{secant}}{\text{adj}}(\theta) = \sec \theta = \frac{\text{hyp}}{\text{adj}}$

 $\frac{\text{cotangent}}{(\theta)} = \cot \theta = \frac{\operatorname{adj}}{\operatorname{opp}}$

StudyTip

 $\tan \theta = \frac{opp}{adi}$



Example 1 Find Values of Trigonometric Ratios

Find the exact values of the six trigonometric functions of θ .

The length of the side opposite θ is 8, the length of the side adjacent to θ is 15, and the length of the hypotenuse is 17.



GuidedPractice



Consider $\sin \theta$ in the figure.

Using $\triangle ABC$: sin $\theta = \frac{BC}{AB}$

Using
$$\triangle AB'C'$$
: sin $\theta = \frac{B'C'}{AB'}$



17

15

Notice that the triangles are similar because they are two right triangles that share a common angle, θ . Because the triangles are similar, the ratios of the corresponding sides are equal. So, $\frac{BC}{AB} = \frac{B'C'}{AB'}$.

Therefore, sin θ has the same value regardless of the triangle used. The values of the functions are constant for a given angle measure. They do not depend on the size of the right triangle.

Example 2 Use One Trigonometric Value to Find Others

If $\cos \theta = \frac{2}{5}$, find the exact values of the five remaining trigonometric functions for the acute angle θ .

Begin by drawing a right triangle and labeling one acute angle θ .

Because $\cos \theta = \frac{\text{adj}}{\text{hyp}} = \frac{2}{5'}$ label the adjacent side 2 and the hypotenuse 5.

By the Pythagorean Theorem, the length of the leg opposite θ is $\sqrt{5^2 - 2^2}$ or $\sqrt{21}$.



GuidedPractice

2. If $\tan \theta = \frac{1}{2'}$ find the exact values of the five remaining trigonometric functions for the acute angle θ .

WatchOut!

Common Misconception

In Example 2, the adjacent side of the triangle could also have been labeled 4 and the hypotenuse 10. This is because $\cos \theta = \frac{2}{5}$ gives the ratio of the adjacent side and hypotenuse, not their specific measures.

You will often be asked to find the trigonometric functions of specific acute angle measures. The table below gives the values of the six trigonometric functions for three common angle measures: 30° , 45° , and 60° . To remember these values, you can use the properties of 30° - 60° - 90° and 45° - 45° - 90° triangles.



You will verify some of these values in Exercises 57-62.

2 Solving Right Triangles Trigonometric functions can be used to find missing side lengths and angle measures of right triangles.

Example 3 Find a Missing Side Length

Find the value of *x*. Round to the nearest tenth, if necessary.

Because you are given an acute angle measure and the length of the hypotenuse of the triangle, use the cosine function to find the length of the side adjacent to the given angle.

$\cos \theta = \frac{\mathrm{adj}}{\mathrm{hyp}}$	Cosine function
$\cos 42^\circ = \frac{x}{18}$	$\theta = 42^\circ$, adj = x, and hyp = 18
$18\cos 42^\circ = x$	Multiply each side by 18.
$13.4 \approx x$	Use a calculator.

Therefore, x is about 13.4.

CHECK You can check your answer by substituting x = 13.4 into $\cos 42^\circ = \frac{x}{18}$.

$$\cos 42^\circ = \frac{x}{18}$$

 $\cos 42^\circ = \frac{13.4}{18}$ $x = 13.4$
 $0.74 = 0.74$ Simplify.

GuidedPractice



TechnologyTip

Degree Mode To evaluate a trigonometric function of an angle measured in degrees, first set the calculator to *degree mode* by selecting DEGREE on the MODE feature of the graphing calculator.





Real-WorldLink

The Ironman Triathlon held in Kailua-Kona Bay, Hawaii, consists of three endurance events, including a 2.4-mile swim, a 112-mile bike ride, and a 26.2-mile marathon.

Source: World Triathlon Corporation

ReadingMath Inverse Trigonometric Ratios The expression $\sin^{-1} x$ is read *the inverse sine of x*. Be careful not to confuse this notation with the notation for negative exponents: $\sin^{-1} x \neq \frac{1}{\sin x}$. Instead, this notation is similar to the notation for an inverse function, $f^{-1}(x)$.

Real-World Example 4 Finding a Missing Side Length

TRIATHLONS A competitor in a triathlon is running along the course shown. Determine the length in feet that the runner must cover to reach the finish line.

An acute angle measure and the opposite side length are given, so the sine function can be used to find the hypotenuse.

$\sin\theta = \frac{\text{opp}}{\text{hyp}}$	Sine function
$\sin 63^\circ = \frac{200}{x}$	$\theta = 63^{\circ}$, opp = 200, and hyp = 2
$x\sin 63^\circ = 200$	Multiply each side by x.
$x = \frac{200}{\sin 63^{\circ}}$ or about 224.47	Divide each side by sin 63°.

63° x ft 200 ft

So, the competitor must run about 224.5 feet to finish the triathlon.

GuidedPractice

4. TRIATHLONS Suppose a competitor in the swimming portion of the race is swimming along the course shown. Find the distance the competitor must swim to reach the shore.



connectED.mcgraw-hill.com

When a trigonometric value of an acute angle is known, the corresponding **inverse trigonometric function** can be used to find the measure of the angle.

1	KeyConcept Inver	se Trigonometric Functions
2	Inverse Sine	If θ is an acute angle and the sine of θ is x, then the inverse sine of x is the measure of angle θ . That is, if sin $\theta = x$, then sin ⁻¹ $x = \theta$.
	Inverse Cosine	If θ is an acute angle and the cosine of θ is x, then the inverse cosine of x is the measure of angle θ . That is, if $\cos \theta = x$, then $\cos^{-1} x = \theta$.
	Inverse Tangent	If θ is an acute angle and the tangent of θ is x , then the inverse tangent of x is the measure of angle θ . That is, if tan $\theta = x$, then $\tan^{-1} x = \theta$.

Example 5 Find a Missing Angle Measure



Douglas Peebles/Alamy

Some applications of trigonometry use an angle of elevation or depression. An **angle of elevation** is the angle formed by a horizontal line and an observer's line of sight to an object above. An **angle of depression** is the angle formed by a horizontal line and an observer's line of sight to an object below.



In the figure, the angles of elevation and depression are congruent because they are alternate interior angles of parallel lines.



AIRPLANES A ground crew worker who is 6 feet tall is directing a plane on a runway. If the worker sights the plane at an angle of elevation of 32°, what is the horizontal distance from the worker to the plane?



Because the worker is 6 feet tall, the vertical distance from the worker to the plane is 150 - 6, or 144 feet. Because the measures of an angle and opposite side are given in the problem, you can use the tangent function to find *x*.

$\tan \theta = \frac{\text{opp}}{\text{adj}}$	Tangent function
$\tan 32^\circ = \frac{144}{x}$	$\theta = 32^{\circ}$, opp = 144, and adj = 2
$x \tan 32^\circ = 144$	Multiply each side by x.
$x = \frac{144}{\tan 32^\circ}$	Divide each side by tan 32°.
$r \approx 230.4$	lise a calculator

So, the horizontal distance from the worker to the plane is approximately 230.4 feet.

GuidedPractice

6. CAMPING A group of hikers on a camping trip climb to the top of a 1500-foot mountain. When the hikers look down at an angle of depression of 36°, they can see the campsite in the distance. What is the horizontal distance between the campsite and the group to the nearest foot?



Real-WorldCareer Airport Ground Crew Ground crewpersons operate ramp-servicing vehicles, handle cargo/baggage, and marshal or

tow aircraft. Crewpersons must have a high school diploma, a valid driver's license, and a good driving record. Angles of elevation or depression can be used to estimate the distance between two objects, as well as the height of an object when given two angles from two different positions of observation.

StudyTip

Indirect Measurement When calculating the distance between two objects using angles of depression, it is important to remember that the objects must lie in the same horizontal plane.

Real-World Example 7 Use Two Angles of Elevation or Depression

BALLOONING A hot air balloon that is moving above a neighborhood has an angle of depression of 28° to one house and 52° to a house down the street. If the height of the balloon is 650 feet, estimate the distance between the two houses.

28°

x ft

X

650 ft

52°

y ft

Draw a diagram to model this situation. Because the angle of elevation from a house to the balloon is congruent to the angle of depression from the balloon to that house, you can label the angles of elevation as shown. Label the horizontal distance from the balloon to the first house *x* and the distance between the two houses *y*.

From the smaller right triangle, you can use the tangent function to find *x*.

$\tan \theta = \frac{\operatorname{opp}}{\operatorname{adj}}$	Tangent function
$\tan 52^\circ = \frac{650}{x}$	$\theta = 52^{\circ}$, opp = 650, and adj =
$x \tan 52^\circ = 650$	Multiply each side by x.
$x = \frac{650}{\tan 52^\circ}$	Divide each side by tan 52°.

From the larger triangle, you can use the tangent function to find *y*.

$\tan \theta = \frac{\text{opp}}{\text{adj}}$	Tangent function
$\tan 28^\circ = \frac{650}{x+y}$	$\theta = 28^\circ$, opp = 650, and adj = $x + y$
$(x+y)\tan 28^\circ = 650$	Multiply each side by $x + y$.
$x + y = \frac{650}{\tan 28^\circ}$	Divide each side by tan 28°.
$\frac{650}{\tan 52^{\circ}} + y = \frac{650}{\tan 28^{\circ}}$	Substitute $x = \frac{650}{\tan 52^\circ}$.
$y = \frac{650}{\tan 28^{\circ}} - \frac{650}{\tan 52^{\circ}}$	Subtract $\frac{650}{\tan 52^{\circ}}$ from each side.

TechnologyTip Using Parentheses When

evaluating a trigonometric expression using a graphing calculator, be careful to close parentheses. While a calculator returns the same value for the expressions tan(26 and tan(26), it does not for expressions tan(26 + 50 and tan(26) + 50.

Therefore, the houses are about 714.6 feet apart.

 $y \approx 714.6$

GuidedPractice

7. BUILDINGS The angle of elevation from a car to the top of an apartment building is 48°. If the angle of elevation from another car that is 22 feet directly in front of the first car is 64°, how tall is the building?

Use a calculator.



Trigonometric functions and inverse relations can be used to **solve a right triangle**, which means to find the measures of all of the sides and angles of the triangle.

ReadingMath

Labeling Triangles

Throughout this chapter, a capital letter will be used to represent both a vertex of a triangle and the measure of the angle at that vertex. The same letter in lowercase will be used to represent both the side opposite that angle and the length of that side.

Example 8 Solve a Right Triangle

Solve each triangle. Round side lengths to the nearest tenth and angle measures to the nearest degree.



a.

Find *x* and *z* using trigonometric functions.

$\tan 35^\circ = \frac{x}{10}$	Substitute.	$\cos 35^\circ = \frac{10}{z}$	Substitute.
10 tan $35^{\circ} = x$	Multiply.	$z \cos 35^\circ = 10$	Multiply.
$7.0 \approx x$	Use a calculator.	$z = \frac{10}{\cos 35^{\circ}}$	Divide.
		$z \approx 12.2$	Use a calculator.

Because the measures of two angles are given, Y can be found by subtracting X from 90°.

$35^\circ + Y = 90^\circ$	Angles X and Y are complementary	
$Y = 55^{\circ}$	Subtract.	

Therefore, $Y = 55^\circ$, $x \approx 7.0$, and $z \approx 12.2$.



Because two side lengths are given, you can use the Pythagorean Theorem to find that $k = \sqrt{185}$ or about 13.6. You can find *J* by using any of the trigonometric functions.

tan
$$J = \frac{11}{8}$$

 $J = \tan^{-1} \frac{11}{8}$
 $J \approx 53.97^{\circ}$ Use a calculator.

Because *J* is now known, you can find *L* by subtracting *J* from 90°.

$$53.97^{\circ} + L \approx 90^{\circ}$$
 Angles J and L are complementary.

 $L \approx 36.03^{\circ}$ Subtract.

Therefore, $J \approx 54^\circ$, $L \approx 36^\circ$, and $k \approx 13.6$.

GuidedPractice



Exercises



Find the exact values of the six trigonometric functions of θ .

Use the given trigonometric function value of the acute angle θ to find the exact values of the five remaining trigonometric function values of θ . (Example 2)

9.	$\sin\theta = \frac{4}{5}$	10.	$\cos \theta = \frac{6}{7}$
11.	$\tan\theta=3$	12.	$\sec \theta = 8$
13.	$\cos\theta = \frac{5}{9}$	14.	$\tan \theta = \frac{1}{4}$
15.	$\cot \theta = 5$	16.	$\csc \theta = 6$
17.	$\sec \theta = \frac{9}{2}$	18.	$\sin \theta = \frac{8}{13}$

Find the value of *x*. Round to the nearest tenth, if necessary. (Example 3)



27 MOUNTAIN CLIMBING A team of climbers must determine the width of a ravine in order to set up equipment to cross it. If the climbers walk 25 feet along the ravine from their crossing point, and sight the crossing point on the far side of the ravine to be at a 35° angle, how wide is the ravine? (Example 4)



- **28. SNOWBOARDING** Brad built a snowboarding ramp with a height of 3.5 feet and an 18° incline. (Example 4)
 - a. Draw a diagram to represent the situation.
 - **b.** Determine the length of the ramp.
- **29. DETOUR** Traffic is detoured from Elwood Ave., left 0.8 mile on Maple St., and then right on Oak St., which intersects Elwood Ave. at a 32° angle. (Example 4)
 - **a.** Draw a diagram to represent the situation.
 - **b.** Determine the length of Elwood Ave. that is detoured.
- **30. PARACHUTING** A paratrooper encounters stronger winds than anticipated while parachuting from 1350 feet, causing him to drift at an 8° angle. How far from the drop zone will the paratrooper land? (Example 4)



Find the measure of angle θ . Round to the nearest degree, if necessary. (Example 5)



39. PARASAILING Kayla decided to try parasailing. She was strapped into a parachute towed by a boat. An 800-foot line connected her parachute to the boat, which was at a 32° angle of depression below her. How high above the water was Kayla? (Example 6)



40. OBSERVATION WHEEL The London Eye is a 135-meter-tall observation wheel. If a passenger at the top of the wheel sights the London Aquarium at a 58° angle of depression, what is the distance between the aquarium and the London Eye? (Example 6)



- **41. ROLLER COASTER** On a roller coaster, 375 feet of track ascend at a 55° angle of elevation to the top before the first and highest drop. (Example 6)
 - a. Draw a diagram to represent the situation.
 - **b.** Determine the height of the roller coaster.
- **42. SKI LIFT** A company is installing a new ski lift on a 225-meter-high mountain that will ascend at a 48° angle of elevation. (Example 6)
 - **a.** Draw a diagram to represent the situation.
 - **b.** Determine the length of cable the lift requires to extend from the base to the peak of the mountain.
- **43. BASKETBALL** Both Derek and Sam are 5 feet 10 inches tall. Derek looks at a 10-foot basketball goal with an angle of elevation of 29°, and Sam looks at the goal with an angle of elevation of 43°. If Sam is directly in front of Derek, how far apart are the boys standing? (Example 7)



- **44. PARIS** A tourist on the first observation level of the Eiffel Tower sights the Musée D'Orsay at a 1.4° angle of depression. A tourist on the third observation level, located 219 meters directly above the first, sights the Musée D'Orsay at a 6.8° angle of depression. (Example 7)
 - a. Draw a diagram to represent the situation.
 - **b.** Determine the distance between the Eiffel Tower and the Musée D'Orsay.

- 45 LIGHTHOUSE Two ships are spotted from the top of a 156-foot lighthouse. The first ship is at a 27° angle of depression, and the second ship is directly behind the first at a 7° angle of depression. (Example 7)
 - a. Draw a diagram to represent the situation.
 - **b.** Determine the distance between the two ships.
- **46. MOUNT RUSHMORE** The faces of the presidents at Mount Rushmore are 60 feet tall. A visitor sees the top of George Washington's head at a 48° angle of elevation and his chin at a 44.76° angle of elevation. Find the height of Mount Rushmore. (Example 7)



Solve each triangle. Round side lengths to the nearest tenth and angle measures to the nearest degree. (Example 8)



- **55. BASEBALL** Michael's seat at a game is 65 feet behind home plate. His line of vision is 10 feet above the field.
 - **a.** Draw a diagram to represent the situation.
 - **b.** What is the angle of depression to home plate?
- **56. HIKING** Jessica is standing 2 miles from the center of the base of Pikes Peak and looking at the summit of the mountain, which is 1.4 miles from the base.
 - a. Draw a diagram to represent the situation.
 - **b.** With what angle of elevation is Jessica looking at the summit of the mountain?

Find the exact value of each expression without using a calculator.

57.	sin 60°	58.	$\cot 30^{\circ}$	59.	sec 30°
60.	$\cos 45^{\circ}$	61.	tan 60°	62.	csc 45°

Without using a calculator, find the measure of the acute angle θ in a right triangle that satisfies each equation.

63. $\tan \theta = 1$ **64.** $\cos \theta = \frac{\sqrt{3}}{2}$ **65.** $\cot \theta = \frac{\sqrt{3}}{3}$ **66.** $\sin \theta = \frac{\sqrt{2}}{2}$ **67.** $\csc \theta = 2$ **68.** $\sec \theta = 2$

Without using a calculator, determine the value of *x*.



71. SCUBA DIVING A scuba diver located 20 feet below the surface of the water spots a shipwreck at a 70° angle of depression. After descending to a point 45 feet above the ocean floor, the diver sees the shipwreck at a 57° angle of depression. Draw a diagram to represent the situation, and determine the depth of the shipwreck.

Find the value of $\cos \theta$ if θ is the measure of the smallest angle in each type of right triangle.

- **72.** 3-4-5 **73.** 5-12-13
- **74. SOLAR POWER** Find the total area of the solar panel shown below.



Without using a calculator, insert the appropriate symbol >, <, or = to complete each equation.

75.	sin 45°	cot 60°	76.	tan 60°	cot 30°
77.	cos 30°	$\csc 45^{\circ}$	78.	cos 30°	$\sin 60^{\circ}$
79.	sec 45°	csc 60°	80.	tan 45°	sec 30°

81. ENGINEERING Determine the depth of the shaft at the large end d of the air duct shown below if the taper of the duct is 3.5° .



- **82. WULTIPLE REPRESENTATIONS** In this problem, you will investigate trigonometric functions of acute angles and their relationship to points on the coordinate plane.
 - **a. GRAPHICAL** Let P(x, y) be a point in the first quadrant. Graph the line through point *P* and the origin. Form a right triangle by connecting the points *P*, (*x*, 0), and the origin. Label the lengths of the legs of the triangle in terms of *x* or *y*. Label the length of the hypotenuse as *r* and the angle the line makes with the *x*-axis θ .
 - **b. ANALYTICAL** Express the value of *r* in terms of *x* and *y*.
 - **c. ANALYTICAL** Express $\sin \theta$, $\cos \theta$, and $\tan \theta$ in terms of *x*, *y*, and/or *r*.
 - **d. VERBAL** Under what condition can the coordinates of point *P* be expressed as $(\cos \theta, \sin \theta)$?
 - **e. ANALYTICAL** Which trigonometric ratio involving θ corresponds to the slope of the line?
 - **f. ANALYTICAL** Find an expression for the slope of the line perpendicular to the line in part **a** in terms of *θ*.

H.O.T. Problems Use Higher-Order Thinking Skills

- **83. PROOF** Prove that if θ is an acute angle of a right triangle, then $\tan \theta = \frac{\sin \theta}{\cos \theta}$ and $\cot \theta = \frac{\cos \theta}{\sin \theta}$.
- **84. ERROR ANALYSIS** Jason and Nadina know the value of $\sin \theta = a$ and are asked to find $\csc \theta$. Jason says that this is not possible, but Nadina disagrees. Is either of them correct? Explain your reasoning.
- **85.** WRITING IN MATH Explain why the six trigonometric functions are transcendental functions.
- **86. CHALLENGE** Write an expression in terms of θ for the area of the scalene triangle shown. Explain.



PROOF Prove that if θ is an acute angle of a right triangle, then $(\sin \theta)^2 + (\cos \theta)^2 = 1$.

REASONING If *A* and *B* are the acute angles of a right triangle and $m \angle A < m \angle B$, determine whether each statement is true or false. If false, give a counterexample.

- **88.** $\sin A < \sin B$
- **89.** $\cos A < \cos B$
- **90.** tan *A* < tan *B*
- **91.** WRITING IN MATH Notice on a graphing calculator that there is no key for finding the secant, cosecant, or cotangent of an angle measure. Explain why you think this might be so.

Spiral Review

92. ECONOMICS The Consumer Price Index (CPI) measures inflation. It is based on the average prices of goods and services in the United States, with the annual average for the years 1982–1984 set at an index of 100. The table shown gives some annual average CPI values from 1955 to 2005. Find an exponential model relating this data (year, CPI) by linearizing the data. Let x = 0 represent 1955. Then use your model to predict the CPI for 2025. (Lesson 3-5)

Solve each equation. Round to the nearest hundredth. (Lesson 3-4)

93. $e^{5x} = 24$ **94.** $2e^{x-7} - 6 = 0$

Sketch and analyze the graph of each function. Describe its domain, range, intercepts, asymptotes, end behavior, and where the function is increasing or decreasing. (Lesson 3-1)

95. $f(x) = -3^{x-2}$ **96.** $f(x) = 2^{3x-4} + 1$ **97.** $f(x) = -4^{-x+6}$

Solve each equation. (Lesson 2-5)

98.
$$\frac{x^2 - 16}{(x+4)(2x-1)} = \frac{4}{x+4} - \frac{1}{2x-1}$$
 99. $\frac{x^2 - 7}{(x+1)(x-5)} = \frac{6}{x+1} + \frac{3}{x-5}$ **100.** $\frac{2x^2 + 3}{3x^2 + 5x + 2} = \frac{5}{3x+2} - \frac{1}{x+1}$

101. NEWSPAPERS The circulation in thousands of papers of a national newspaper is shown. (Lesson 2-1)

Year	2002	2003	2004	2005	2006	2007	2008
Circulation (in thousands)	904.3	814.7	773.9	725.5	716.2	699.1	673.0

a. Let *x* equal the number of years after 2001. Create a scatter plot of the data.

b. Determine a power function to model the data.

c. Use the function to predict the circulation of the newspaper in 2015.

Skills Review for Standardized Tests



104. A person holds one end of a rope that runs through a pulley and has a weight attached to the other end. Assume that the weight is at the same height as the person's hand. What is the distance from the person's hand to the weight?



- A 7.8 ft
- **B** 10.5 ft
- C 12.9 ft
- **D** 14.3 ft
- **105. REVIEW** A kite is being flown at a 45° angle. The string of the kite is 120 feet long. How high is the kite above the point at which the string is held?
 - **F** 60 ft
 - G $60\sqrt{2}$ ft
 - H $60\sqrt{3}$ ft
 - J 120 ft

Year	CPI
1955	26.8
1965	31.5
1975	53.8
1985	107.6
1995	152.4
2005	195.3

Source: Bureau of Labor Statistics

Degrees and Radians

E		rees and ka	idians 🗠	P 0
Then	Now	Why?		
• You used the measures of acute angles in triangles given in degrees. (Lesson 4-1)	 Convert degree measures of ang to radian measure and vice versa. Use angle measure to solve real-wor problems. 	 In Lesson 4-1, you wo angles, but angles can measurement. For exa a 540 is an aerial trick and the board rotate th or one and a half complete 	rked only with acute have <i>any</i> real number imple, in skateboarding, in which a skateboarder hrough an angle of 540°, plete turns, in midair.	
NewVocabu vertex initial side terminal side standard position radian	lary 1 Angles two none thought of as perspective, t position after vertex at the	and Their Measures Fro ollinear rays that share a cor being formed by the action of the starting position of the ra rotation forms the angle's te origin and its initial side alor	om geometry, you may recall a mmon endpoint known as a v of rotating a ray about its end ay forms the initial side of the erminal side . In the coordinate ng the positive x-axis is said to	an angle being defined as ertex. An angle can also be point. From this dynamic angle, while the ray's e plane, an angle with its o be in <mark>standard position</mark> .
coterminal angles linear speed angular speed sector		Angle terminal side	Angle in Standard Post	sition

The measure of an angle describes the amount and direction of rotation necessary to move from the initial side to the terminal side of the angle. A positive angle is generated by a counterclockwise rotation and a *negative angle* by a clockwise rotation.



initial side

vertex

The most common angular unit of measure is the *degree* (°), which is equivalent to $\frac{1}{360}$ of a full rotation (counterclockwise) about the vertex. From the diagram shown, you can see that 360° corresponds to 1 complete rotation, 180° to a $\frac{1}{2}$ rotation, 90° to a $\frac{1}{4}$ rotation, and so on, as marked along the circumference of the circle.



23

initial side

StudyTip

Base 60 The concept of degree measurement dates back to the ancient Babylonians, who made early astronomical calculations using their number system, which was based on 60 (sexagesimal) rather than on 10 (decimal) as we do today.

TechnologyTip

DMS Form You can use some calculators to convert decimal degree values to degrees, minutes, and seconds using the DMS function under the Angle menu. Degree measures can also be expressed using a decimal degree form or a degree-minute-second (DMS) form where each degree is subdivided into 60 minutes (') and each minute is subdivided into 60 seconds (").

Example 1 Convert Between DMS and Decimal Degree Form

Write each decimal degree measure in DMS form and each DMS measure in decimal degree form to the nearest thousandth.

```
a. 56.735°
```

First, convert 0.735° into minutes and seconds.

$$56.735^{\circ} = 56^{\circ} + 0.735^{\circ} \left(\frac{60'}{1^{\circ}}\right) \qquad 1^{\circ} = 60^{\circ}$$
$$= 56^{\circ} + 44.1' \qquad \text{Simplify}$$

Next, convert 0.1' into seconds.

$$56.735^{\circ} = 56^{\circ} + 44' + 0.1' \left(\frac{60''}{1'}\right) \qquad 1' = 60'$$

= 56^{\circ} + 44' + 6'' Simplify

Therefore, 56.735° can be written as $56^{\circ} 44' 6''$.

b. 32° 5′ 28″

Each minute is $\frac{1}{60}$ of a degree and each second is $\frac{1}{60}$ of a minute, so each second is $\frac{1}{3600}$ of a degree.

 $32^{\circ} 5' 28'' = 32^{\circ} + 5' \left(\frac{1^{\circ}}{60'}\right) + 28'' \left(\frac{1^{\circ}}{3600''}\right) \qquad 1' = \frac{1}{60} (1^{\circ}) \text{ and } 1'' = \frac{1}{3600} (1^{\circ})$ $\approx 32^{\circ} + 0.083 + 0.008 \qquad \text{Simplify.}$ $\approx 32.091^{\circ} \qquad \text{Add.}$

Therefore, 32° 5′ 28″ can be written as about 32.091°.

GuidedPractice

1A. 213.875°

1B. 89° 56′ 7″

Measuring angles in degrees is appropriate when applying trigonometry to solve many real-world problems, such as those in surveying and navigation. For other applications with trigonometric functions, using an angle measured in degrees poses a significant problem. A degree has no relationship to any linear measure; inch-degrees or $\frac{inch}{degree}$ has no meaning. Measuring angles in radians provides a solution to this problem.

Key Concept	Radian Measure	
Words	The measure θ in radians of a central angle of a circle is equal to the ratio of the length of the intercepted arc s to the radius r of the circle.	() () () () () () () () () () () () () (
Symbols	$\theta = \frac{s}{r}$, where θ is measured in radians (rad)	
Example	A central angle has a measure of 1 radian if it intercepts an arc with the same length as the radius of the circle.	$\theta = 1$ radian when $s = r$.

Notice that as long as the arc length *s* and radius *r* are measured using the same linear units, the ratio $\frac{s}{r}$ is unitless. For this reason, the word *radian* or its abbreviation *rad* is usually omitted when writing the radian measure of an angle.

StudyTip

Degree-Radian Equivalences From the equivalence statement shown, you can determine that $1^{\circ} \approx 0.017$ rad and $1 \text{ rad} \approx 57.296^{\circ}$. The central angle representing one full rotation counterclockwise about a vertex corresponds to an arc length equivalent to the circumference of the circle, $2\pi r$. From this, you can obtain the following radian measures.



Because 2π radians and 360° both correspond to one complete revolution, you can write $360^{\circ} = 2\pi$ radians or $180^{\circ} = \pi$ radians. This last equation leads to the following equivalence statements.

$$1^{\circ} = \frac{\pi}{180}$$
 radians and $1 \text{ radian} = \left(\frac{180}{\pi}\right)$

Using these statements, we obtain the following conversion rules.

KeyConcept Degree/Radian Conversion Rules

- 1. To convert a degree measure to radians, multiply by $\frac{\pi \text{ radians}}{190^\circ}$
- 2. To convert a radian measure to degrees, multiply by $\frac{180^{\circ}}{\pi \text{ radians}}$

Example 2 Convert Between Degree and Radian Measure

Write each degree measure in radians as a multiple of π and each radian measure in degrees.

a. 120°
120° = 120°
$$\left(\frac{\pi \operatorname{radians}}{180°}\right)$$
 Multiply by $\frac{\pi \operatorname{radians}}{180°}$.
 $= \frac{2\pi}{3} \operatorname{radians} \operatorname{or} \frac{2\pi}{3}$ Simplify.
b. -45°
 $-45° = -45° \left(\frac{\pi \operatorname{radians}}{180°}\right)$ Multiply by $\frac{\pi \operatorname{radians}}{180°}$.
 $= -\frac{\pi}{4} \operatorname{radians} \operatorname{or} -\frac{\pi}{4}$ Simplify.
c. $\frac{5\pi}{6}$
 $\frac{5\pi}{6} = \frac{5\pi}{6} \operatorname{radians}$ Multiply by $\frac{180°}{\pi \operatorname{radians}}$.
 $= \frac{5\pi}{6} \operatorname{radians} \left(\frac{180°}{\pi \operatorname{radians}}\right) \operatorname{or} 150°$ Simplify.
d. $-\frac{3\pi}{2}$
 $-\frac{3\pi}{2} = -\frac{3\pi}{2} \operatorname{radians} \left(\frac{180°}{\pi \operatorname{radians}}\right) \operatorname{or} -270°$ Simplify.
EquidedPractice
24. $210°$ 28. $-60°$ 2C. $\frac{4\pi}{3}$ 2D. $-\frac{\pi}{6}$

connectED.mcgraw-hill.com

233

ReadingMath

Angle Measure If no units of angle measure are specified, radian measure is implied. If degrees are intended, the degree symbol (°) must be used.

ReadingMath

Naming Angles In trigonometry, angles are often labeled using Greek letters, such as α (alpha), β (beta), and θ (theta).

By defining angles in terms of their rotation about a vertex, two angles can have the same initial and terminal sides but different measures. Such angles are called coterminal angles. In the figures below, angles α and β are coterminal.





The two positive coterminal angles shown differ by one full rotation. A given angle has infinitely many coterminal angles found by adding or subtracting integer multiples of 360° or 2π radians.

KeyConcept Coterminal Angles		
Degrees	Radians	
If α is the degree measure of an angle, then all angles measuring α + 360 n° , where n is an integer, are coterminal with α .	If α is the radian measure of an angle, then all angles measuring $\alpha + 2n\pi$, where <i>n</i> is an integer, are coterminal with α .	

Example 3 Find and Draw Coterminal Angles

Identify all angles that are coterminal with the given angle. Then find and draw one positive and one negative angle coterminal with the given angle.





2 Applications with Angle Measure Solving $\theta = \frac{s}{r}$ for the arc length *s* yields a convenient formula for finding the length of an arc of a circle.



When θ is measured in degrees, you could also use the equation $s = \frac{\pi r \theta}{180}$, which already incorporates the degree-radian conversion.

Example 4 Find Arc Length

Find the length of the intercepted arc in each circle with the given central angle measure and radius. Round to the nearest tenth.



s =

The length of the intercepted arc is $\frac{5\pi}{4}$ or about 3.9 centimeters.

b. $60^{\circ}, r = 2$ in. Method 1



$$60^{\circ} = 60^{\circ} \left(\frac{\pi \text{ radians}}{180^{\circ}}\right) \qquad \text{Muttiply by } \frac{\pi \text{ radians}}{180^{\circ}}$$
$$= \frac{\pi}{3} \qquad \text{Simplify.}$$

Substitute r = 2 and $\theta = \frac{\pi}{3}$.

$$r\theta$$
 Arc length
 $2\left(\frac{\pi}{3}\right)$ $r=2$ and $\theta=\frac{\pi}{3}$

Simplify.

2 and $\theta = 60^{\circ}$

$$=\frac{2\pi}{3}$$

Method 2 Use
$$s = \frac{\pi r \theta}{180^{\circ}}$$
 to find the arc length.
 $s = \frac{\pi r \theta}{180^{\circ}}$ Arc length

$$= \frac{\pi(2)(60^\circ)}{180^\circ} \qquad r = 2 \text{ an}$$
$$= \frac{2\pi}{3} \qquad \text{Simplify.}$$

The length of the intercepted arc is $\frac{2\pi}{3}$ or about 2.1 inches.

GuidedPractice

4A.
$$\frac{2\pi}{3}$$
, $r = 2$ m

4B. 135°, *r* = 0.5 ft



Operating with Radians Notice in Example 4a that when r = 5 centimeters and $\theta = \frac{\pi}{4}$ radians, $s = \frac{5\pi}{4}$ centimeters, not $\frac{5\pi}{4}$ centimeter-radians. This is because a radian is a unitless ratio.

60°

2 in

The formula for arc length can be used to analyze circular motion. The rate at which an object moves along a circular path is called its **linear speed**. The rate at which the object *rotates* about a fixed point is called its **angular speed**. Linear speed is measured in units like miles per hour, while angular speed is measured in units like revolutions per minute.

KeyConcept Linear and Angular Speed

Suppose an object moves at a constant speed along a circular path of radius r.

If *s* is the arc length traveled by the object during time *t*, then the object's *linear* speed *v* is given by $v = \frac{S}{4}$.

If θ is the angle of rotation (in radians) through which the object moves during time *t*, then the *angular speed* ω of the object is given by $\omega = \frac{\theta}{t}$.



Real-World Example 5 Find Angular and Linear Speeds

- **BICYCLING** A bicycle messenger rides the bicycle shown.
- **a**. During one delivery, the tires rotate at a rate of 140 revolutions per minute. Find the angular speed of the tire in radians per minute.

Because each rotation measures 2π radians, 140 revolutions correspond to an angle of rotation θ of 140 × 2π or 280 π radians.

$$\omega = \frac{\theta}{t}$$
Angular speed
$$= \frac{280\pi \text{ radians}}{1 \text{ minute}}$$
 $\theta = 280\pi \text{ radians and } t =$



Therefore, the angular speed of the tire is 280π or about 879.6 radians per minute.

b. On part of the trip to the next delivery, the tire turns at a constant rate of 2.5 revolutions per second. Find the linear speed of the tire in miles per hour.

1 minute

A rotation of 2.5 revolutions corresponds to an angle of rotation θ of 2.5 × 2 π or 5 π .

 $v = \frac{s}{t}$ Linear speed $= \frac{r\theta}{t}$ $= \frac{15(5\pi) \text{ inches}}{1 \text{ second}} \text{ or } \frac{75\pi \text{ inches}}{1 \text{ second}}$ $r = 15 \text{ inches}, \theta = 5\pi \text{ radians, and } t = 1 \text{ second}$

Use dimensional analysis to convert this speed from inches per second to miles per hour.



Therefore, the linear speed of the tire is about 13.4 miles per hour.

GuidedPractice

MEDIA Consider the DVD shown.

- **5A.** Find the angular speed of the DVD in radians per second if the disc rotates at a rate of 3.5 revolutions per second.
- **5B.** If the DVD player overheats and the disc begins to rotate at a slower rate of 3 revolutions per second, find the disc's linear speed in meters per minute.



ReadingMath

Omega The lowercase Greek letter omega ω is usually used to denote angular speed.



Real-WorldLink

In some U.S. cities, it is possible for bicycle messengers to ride an average of 30 to 35 miles a day while making 30 to 45 deliveries.

Source: New York Bicycle Messenger Association Recall from geometry that a sector of a circle is a region bounded by a central angle and its intercepted arc. For example, the shaded portion in the figure is a sector of circle P. The ratio of the area of a sector to the area of a whole circle is equal to the ratio of the corresponding arc length to the circumference of the circle. Let A represent the area of the sector.

$$\frac{A}{\pi r^2} = \frac{\text{length of } \widehat{QRS}}{2\pi r} \qquad \frac{\text{area of sector}}{\text{area of circle}} = \frac{\text{arc length}}{\text{circumference of circle}}$$

$$\frac{A}{\pi r^2} = \frac{r\theta}{2\pi r} \qquad \text{The length of } \widehat{QRS} \text{ is } r\theta.$$

$$A = \frac{1}{2}r^2\theta \qquad \text{Solve for } A.$$

KeyConcept Area of a Sector

The area A of a sector of a circle with radius r and central angle θ is

$$A = \frac{1}{2}r^2\theta$$



where θ is measured in radians.

Example 6 Find Areas of Sectors

a. Find the area of the sector of the circle. The measure of the sector's central angle θ is $\frac{7\pi}{8}$, and the radius is 3 centimeters.

$$A = \frac{1}{2}r^{2}\theta \qquad \text{Area of a sector}$$
$$= \frac{1}{2}(3)^{2}\left(\frac{7\pi}{8}\right) \text{ or } \frac{63\pi}{14} \qquad r = 3 \text{ and } \theta = \frac{7}{14}$$

3 cm

16 in



130°

b. WIPERS Find the approximate area swept by the wiper blade shown, if the total length of

The area swept by the wiper blade is the difference between the areas of the sectors with radii 26 inches and 26 - 16 or 10 inches.

the windshield wiper mechanism is 26 inches.

Convert the central angle measure to radians.

$$130^\circ = 130^\circ \left(\frac{\pi \text{ radians}}{180^\circ}\right) = \frac{13\pi}{18}$$



$$A = A_1 - A_2$$

Swept area
$$= \frac{1}{2}(26)^2 \left(\frac{13\pi}{18}\right) - \frac{1}{2}(10)^2 \left(\frac{13\pi}{18}\right)$$

Area of a sector
$$= \frac{2197\pi}{9} - \frac{325\pi}{9}$$

Simplify.
$$= 208\pi \text{ or about } 653.5$$

Simplify.

Therefore, the swept area is about 653.5 square inches.

GuidedPractice

Find the area of the sector of a circle with the given central angle θ and radius *r*.

6A. $\theta = \frac{3\pi}{4}, r = 1.5 \text{ ft}$

6B. $\theta = 50^{\circ}, r = 6 \text{ m}$



Ovolker Moehrke/Corbis

Real-WorldLink

A typical wipe angle for a front windshield wiper of a passenger car is about 67°. Windshield wiper blades are generally 12-30 inches long. Source: Car and Driver

Exercises



Write each decimal degree measure in DMS form and each DMS measure in decimal degree form to the nearest thousandth. (Example 1)

1.	11.773°	2.	58.244°
3.	141.549°	4.	273.396°
5.	87° 53′ 10″	6.	126° 6′ 34″
7.	45° 21' 25″	8.	301° 42′ 8″

9. NAVIGATION A sailing enthusiast uses a sextant, an instrument that can measure the angle between two objects with a precision to the nearest 10 seconds, to measure the angle between his sailboat and a lighthouse. If his reading is 17° 37′ 50″, what is the measure in decimal degree form to the nearest hundredth? (Example 1)



Write each degree measure in radians as a multiple of π and each radian measure in degrees. (Example 2)

10.	30°	11.	225°
12.	-165°	13.	-45°

14.
$$\frac{2\pi}{3}$$
 15. $\frac{5\pi}{2}$

16.
$$-\frac{\pi}{4}$$
 17. $-\frac{7\pi}{6}$

Identify all angles that are coterminal with the given angle. Then find and draw one positive and one negative angle coterminal with the given angle. (Example 3)

18.	120°	19.	-75°
20.	225°	21.	-150°
22.	$\frac{\pi}{3}$	23.	$-\frac{3\pi}{4}$
24.	$-\frac{\pi}{12}$	25.	$\frac{3\pi}{2}$

26. GAME SHOW Sofia is spinning a wheel on a game show. There are 20 values in equal-sized spaces around the circumference of the wheel. The value that Sofia needs to win is two spaces above the space where she starts her spin, and the wheel must make at least one full rotation for the spin to count. Describe a spin rotation in degrees that will give Sofia a winning result. (Example 3)



Find the length of the intercepted arc with the given central angle measure in a circle with the given radius. Round to the nearest tenth. (Example 4)

27. $\frac{\pi}{6}$, $r = 2.5$ m	28. $\frac{2\pi}{3}$, $r = 3$ in.
29. $\frac{5\pi}{12}$, $r = 4$ yd	30. 105°, <i>r</i> = 18.2 cm
31. 45°, <i>r</i> = 5 mi	32. 150°, <i>r</i> = 79 mm

33 AMUSEMENT PARK A carousel at an amusement park rotates 3024° per ride. (Example 4)

- **a.** How far would a rider seated 13 feet from the center of the carousel travel during the ride?
- **b.** How much farther would a second rider seated 18 feet from the center of the carousel travel during the ride than the rider in part **a**?

Find the rotation in revolutions per minute given the angular speed and the radius given the linear speed and the rate of rotation. (Example 5)

37. *v* = 82.3 m/s, 131 rev/min

- **34.** $\omega = \frac{2}{3}\pi \text{ rad/s}$ **35.** $\omega = 135\pi \text{ rad/h}$
- **36.** $\omega = 104\pi \, \mathrm{rad}/\mathrm{min}$
- **38.** v = 144.2 ft/min, 10.9 rev/min
- **39.** v = 553 in./h, 0.09 rev/min
- **40. MANUFACTURING** A company manufactures several circular saws with the blade diameters and motor speeds shown below. (Example 5)

Blade Diameter (in.)	Motor Speed (rps)
3	2800
5	5500
$5\frac{1}{2}$	4500
$6\frac{1}{8}$	5500
$7\frac{1}{4}$	5000

- **a.** Determine the angular and linear speeds of the blades in each saw. Round to the nearest tenth.
- **b.** How much faster is the linear speed of the $6\frac{1}{8}$ -inch saw compared to the 3-inch saw?
- **41. CARS** On a stretch of interstate, a vehicle's tires range between 646 and 840 revolutions per minute. The diameter of each tire is 26 inches. (Example 5)
 - **a.** Find the range of values for the angular speeds of the tires in radians per minute.
 - **b.** Find the range of values for the linear speeds of the tires in miles per hour.

42. TIME A wall clock has a face diameter of $8\frac{1}{2}$ inches. The length of the hour hand is 2.4 inches, the length of the minute hand is 3.2 inches, and the length of the second hand is 3.4 inches. (Example 5)



- **a.** Determine the angular speed in radians per hour and the linear speed in inches per hour for each hand.
- **b.** If the linear speed of the second hand is 20 inches per minute, is the clock running fast or slow? How much time would it gain or lose per day?

GEOMETRY Find the area of each sector. (Example 6)



49. GAMES The dart board shown is divided into twenty equal sectors. If the diameter of the board is 18 inches, what area of the board does each sector cover? (Example 6)



50. LAWN CARE A sprinkler waters an area that forms one third of a circle. If the stream from the sprinkler extends 6 feet, what area of the grass does the sprinkler water? (Example 6)

The area of a sector of a circle and the measure of its central angle are given. Find the radius of the circle.

(51) $A = 29 \text{ ft}^2, \theta = 68^\circ$	52. $A = 808 \text{ cm}^2, \theta = 210^\circ$
53. $A = 377 \text{ in}^2, \ \theta = \frac{5\pi}{3}$	54. $A = 75 \text{ m}^2, \ \theta = \frac{3\pi}{4}$

55. Describe the radian measure between 0 and 2π of an angle θ that is in standard position with a terminal side that lies in:

a. Quadrant I	c. Quadrant III
b. Quadrant II	d. Quadrant IV

- **56.** If the terminal side of an angle that is in standard position lies on one of the axes, it is called a *quadrantal angle*. Give the radian measures of four quadrantal angles.
- **57. GEOGRAPHY** Phoenix, Arizona, and Ogden, Utah, are located on the same line of longitude, which means that Ogden is directly north of Phoenix. The latitude of Phoenix is 33° 26' N, and the latitude of Ogden is 41° 12' N. If Earth's radius is approximately 3963 miles, about how far apart are the two cities?



Find the measure of angle θ in radians and degrees.



62. TRACK The curve of a standard 8-lane track is semicircular as shown.



- **a.** What is the length of the outside edge of Lane 4 in the curve?
- **b.** How much longer is the inside edge of Lane 7 than the inside edge of Lane 3 in the curve?



- **63. DRAMA** A pulley with radius *r* is being used to remove part of the set of a play during intermission. The height of the pulley is 12 feet.
 - **a.** If the radius of the pulley is 6 inches and it rotates 180°, how high will the object be lifted?
 - **b.** If the radius of the pulley is 4 inches and it rotates 900°, how high will the object be lifted?



- **64. ENGINEERING** A pulley like the one in Exercise 63 is being used to lift a crate in a warehouse. Determine which of the following scenarios could be used to lift the crate a distance of 15 feet the fastest. Explain how you reached your conclusion.
 - **I.** The radius of the pulley is 5 inches rotating at 65 revolutions per minute.
 - **II.** The radius of the pulley is 4.5 inches rotating at 70 revolutions per minute.
 - **III.** The radius of the pulley is 6 inches rotating at 60 revolutions per minute.

GEOMETRY Find the area of each shaded region.



67. CARS The speedometer shown measures the speed of a car in miles per hour.



- **a.** If the angle between 25 mi/h and 60 mi/h is 81.1°, about how many miles per hour are represented by each degree?
- **b.** If the angle of the speedometer changes by 95°, how much did the speed of the car increase?

Find the complement and supplement of each angle, if possible. If not possible, explain your reasoning.

68. $\frac{2\pi}{5}$ **69.** $\frac{5\pi}{6}$ **70.** $\frac{3\pi}{8}$ **71.** $-\frac{\pi}{3}$

- **72. SKATEBOARDING** A physics class conducted an experiment to test three different wheel sizes on a skateboard with constant angular speed.
 - **a.** Write an equation for the linear speed of the skateboard in terms of the radius and angular speed. Explain your reasoning.
 - **b.** Using the equation you wrote in part **a**, predict the linear speed in meters per second of a skateboard with an angular speed of 3 revolutions per second for wheel diameters of 52, 56, and 60 millimeters.
 - **c.** Based on your results in part **b**, how do you think wheel size affects linear speed?

H.O.T. Problems Use Higher-Order Thinking Skills

- **73. ERROR ANALYSIS** Sarah and Mateo are told that the perimeter of a sector of a circle is 10 times the length of the circle's radius. Sarah thinks that the radian measure of the sector's central angle is 8 radians. Mateo thinks that there is not enough information given to solve the problem. Is either of them correct? Explain your reasoning.
- **74. CHALLENGE** The two circles shown are concentric. If the length of the arc from *A* to *B* measures 8π inches and DB = 2 inches, find the arc length from *C* to *D* in terms of π .



REASONING Describe how the linear speed would change for each parameter below. Explain.

- **75.** a decrease in the radius
- **76.** a decrease in the unit of time
- 77. an increase in the angular speed

78. PROOF If
$$\frac{s_1}{r_1} = \frac{s_2}{r_{2'}}$$
 prove that $\theta_1 = \theta_2$.

- **79. REASONING** What effect does doubling the radius of a circle have on each of the following measures? Explain your reasoning.
 - **a.** the perimeter of the sector of the circle with a central angle that measures *θ* radians
 - **b.** the area of a sector of the circle with a central angle that measures *θ* radians
- **80.** WRITING IN MATH Compare and contrast degree and radian measures. Then create a diagram similar to the one on page 231. Label the diagram using degree measures on the inside and radian measures on the outside of the circle.

Spiral Review

Use the given trigonometric function value of the acute angle θ to find the exact values of the five remaining trigonometric function values of θ . (Lesson 4-1)

81.
$$\sin \theta = \frac{8}{15}$$
 82. $\sec \theta = \frac{4\sqrt{7}}{10}$ **83.** $\cot \theta = \frac{17}{19}$

84. BANKING An account that Hally's grandmother opened in 1955 earned continuously compounded interest. The table shows the balances of the account from 1955 to 1959. (Lesson 3-5)

- **a.** Use regression to find a function that models the amount in the account. Use the number of years after Jan. 1, 1955, as the independent variable.
- **b.** Write the equation from part **a** in terms of base *e*.
- **c.** What was the interest rate on the account if no deposits or withdrawals were made during the period in question?

Express each logarithm in terms of ln 2 and ln 5. (Lesson 3-2)

85.
$$\ln \frac{25}{16}$$
 86. $\ln 250$ **87.** $\ln \frac{10}{25}$

List all possible rational zeros of each function. Then determine which, if any, are zeros. (Lesson 2-4) **88.** $f(x) = x^4 - x^3 - 12x - 144$ **89.** $g(x) = x^3 - 5x^2 - 4x + 20$ **90.** $g(x) = 6x^4 + 35x^3 - x^2 - 7x - 1$ Describe the end behavior of each function. (Lesson 1-3) **91.** $f(x) = 4x^5 + 2x^4 - 3x - 1$ **92.** $g(x) = -x^6 + x^4 - 5x^2 + 4$ **93.** $h(x) = -\frac{1}{x^3} + 2$

Write each set in set-builder and interval notation, if possible. (Lesson 0-1)

94. <i>n</i> > -7	95. −4 ≤ <i>x</i> < 10	96. $y < 1$ or $y \ge 11$
94. $n > -7$	95. $-4 \le x < 10$	96. <i>y</i> < 1 or <i>y</i> ≥ .

Skills Review for Standardized Tests

97. SAT/ACT In the figure, *C* and *D* are the centers of the two circles with radii of 3 and 2, respectively. If the larger shaded region has an area of 9, what is the area of the smaller shaded region?



H 1

J 3

F −1

G 0

99. REVIEW If sec $\theta = \frac{25}{7}$ and θ is acute, then sin $\theta =$

A $\frac{7}{25}$ **B** $\frac{24}{25}$ **C** $-\frac{24}{25}$ **D** $\frac{25}{7}$

(

100. Which of the following radian measures is equal to 56°?

F
$$\frac{\pi}{15}$$
 H $\frac{14\pi}{45}$
G $\frac{7\pi}{45}$ J $\frac{\pi}{3}$

 Jan. 1, 1956
 \$2251.61

 Jan. 1, 1957
 \$2371.79

 Jan. 1, 1958
 \$2498.39

 Jan. 1, 1959
 \$2631.74

Balance

\$2137.52

Date

Jan. 1, 1955

1

2

3

4

5

Trigonometric Functions on the Unit Circle

Now Why? Then You found values Find values of A blood pressure of 120 over 80, measured in millimeters of mercury, of trigonometric trigonometric functions means that a person's blood pressure oscillates or cycles between functions for acute for any angle. 20 millimeters above and below a pressure of 100 millimeters of angles using ratios mercury for a given time t in seconds. A complete cycle of this Find values of oscillation takes about 1 second. in right triangles. trigonometric (Lesson 4-1) functions using the If the pressure exerted by the blood at time t = 0.25 second is 120 millimeters of mercury, then at time t = 1.25 seconds unit circle. the pressure is also 120 millimeters of mercury.

NewVocabulary quadrantal angle reference angle unit circle circular function periodic function period

Bc

Trigonometric Functions of Any Angle In Lesson 4-1, the definitions of the six trigonometric functions were restricted to positive acute angles. In this lesson, these definitions are extended to include *any* angle.



Example 1 Evaluate Trigonometric Functions Given a Point

Let (8, -6) be a point on the terminal side of an angle θ in standard position. Find the exact values of the six trigonometric functions of θ .

Use the values of *x* and *y* to find *r*.

$$r = \sqrt{x^2 + y^2}$$
Pythagorean Theorem $= \sqrt{8^2 + (-6)^2}$ $x = 8$ and $y = -6$ $= \sqrt{100}$ or 10Take the positive square root.



Use x = 8, y = -6, and r = 10 to write the six trigonometric ratios.

$\sin \theta = \frac{y}{r} = \frac{-6}{10} \text{ or } -\frac{3}{5}$	$\cos \theta = \frac{x}{r} = \frac{8}{10} \text{ or } \frac{4}{5}$	$\tan \theta = \frac{y}{x} = \frac{-6}{8} \text{ or } -\frac{3}{4}$
$\csc \theta = \frac{r}{y} = \frac{10}{-6} \text{ or } -\frac{5}{3}$	$\sec \theta = \frac{r}{x} = \frac{10}{8} \text{ or } \frac{5}{4}$	$\cot \theta = \frac{x}{y} = \frac{8}{-6} \text{ or } -\frac{4}{3}$

GuidedPractice

The given point lies on the terminal side of an angle θ in standard position. Find the values of the six trigonometric functions of θ .

1A. (4, 3)

In Example 1, you found the trigonometric values of θ without knowing the measure of θ . Now we will discuss methods for finding these function values when only θ is known. Consider trigonometric functions of quadrantal angles. When the terminal side of an angle θ that is in standard position lies on one of the coordinate axes, the angle is called a **quadrantal angle**.

StudyTip

Quadrantal Angles There are infinitely many quadrantal angles that are coterminal with the quadrantal angles listed at the right. The measure of a quadrantal angle is a multiple of 90° or $\frac{\pi}{2}$.



You can find the values of the trigonometric functions of quadrantal angles by choosing a point on the terminal side of the angle and evaluating the function at that point. Any point can be chosen. However, to simplify calculations, pick a point for which *r* equals 1.

Example 2 Evaluate Trigonometric Functions of Quadrantal Angles

Find the exact value of each trigonometric function, if defined. If not defined, write undefined.

```
a. sin (-180°)
```

The terminal side of -180° in standard position lies on the negative *x*-axis. Choose a point *P* on the terminal side of the angle. A convenient point is (-1, 0) because r = 1.

in
$$(-180^\circ) = \frac{y}{r}$$

= $\frac{0}{1}$ or 0 $y = 0$ and r

$$y = 0$$
 and $r = 1$

b. $\tan \frac{3\pi}{2}$

S

The terminal side of $\frac{3\pi}{2}$ in standard position lies on the negative *y*-axis. Choose a point P(0, -1) on the terminal side of the angle because r = 1.

$$\tan \frac{3\pi}{2} = \frac{y}{x}$$
 Tangent function
= $\frac{-1}{0}$ or undefined $y = -1$ and $x = 0$

c. sec 4π

The terminal side of 4π in standard position lies on the positive *x*-axis. The point (1, 0) is convenient because r = 1.

 $\sec 4\pi = \frac{r}{r}$ Secant function $=\frac{1}{1}$ or 1

r = 1 and x = 1

GuidedPractice

2A. cos 270°

```
2B. \csc \frac{\pi}{2}
```







2C. cot (−90°)

To find the values of the trigonometric functions of angles that are neither acute nor quadrantal, consider the three cases shown below in which *a* and *b* are positive real numbers. Compare the values of sine, cosine, and tangent of θ and θ' .



This angle θ' , called a **reference angle**, can be used to find the trigonometric values of any angle θ .

KeyConcept Reference Angle Rules

If θ is an angle in standard position, its reference angle θ' is the acute angle formed by the terminal side of θ and the *x*-axis. The reference angle θ' for any angle θ , $0^{\circ} < \theta < 360^{\circ}$ or $0 < \theta < 2\pi$, is defined as follows.



To find a reference angle for angles outside the interval $0^\circ < \theta < 360^\circ$ or $0 < \theta < 2\pi$, first find a corresponding coterminal angle in this interval.

Example 3 Find Reference Angles

Sketch each angle. Then find its reference angle.



The terminal side of 300° lies in Quadrant IV. Therefore, its reference angle is $\theta' = 360^\circ - 300^\circ$ or 60° . A coterm termina its reference $\theta = 300^\circ$ $\theta' = 60^\circ$

3B. −240°

b. $-\frac{2\pi}{3}$ A coterminal angle is $2\pi - \frac{2\pi}{3}$ or $\frac{4\pi}{3}$. The terminal side of $\frac{4\pi}{3}$ lies in Quadrant III, so its reference angle is $\frac{4\pi}{3} - \pi$ or $\frac{\pi}{3}$.



3C. 390°

StudyTip Reference Angles Notice that in some cases, the three trigonometric values of θ and θ' (read *theta prime*) are the same. In other cases, they differ only in sign.

GuidedPractice

3A. $\frac{5\pi}{4}$

Because the trigonometric values of an angle and its reference angle are equal or differ only in sign, you can use the following steps to find the value of a trigonometric function of any angle θ .



The signs of the trigonometric functions in each quadrant can be determined using the function definitions given on page 242. For example, because $\sin \theta = \frac{y}{r}$, it follows that $\sin \theta$ is negative when y < 0, which occurs in Quadrants III and IV. Using this same logic, you can verify each of the signs for $\sin \theta$, $\cos \theta$, and $\tan \theta$ shown in the diagram. Notice that these values depend only on *x* and *y* because *r* is always positive.

y				
Quadrant II	Quadrant I			
$\sin \theta$: +	sin θ : +			
$\cos \theta$: –	$\cos \theta$: +			
tan θ: —	tan θ : +			
Quadrant III	Quadrant IV			
sin θ : –	sin θ : –			
$\cos \theta$: –	$\cos \theta$: +			

StudyTip

Memorizing Trigonometric Values To memorize the exact values of sine for 0°, 30°, 45°, 60°, and 90°, consider the following pattern.

$$\sin 0^{\circ} = \frac{\sqrt{0}}{2}, \text{ or } 0$$

$$\sin 30^{\circ} = \frac{\sqrt{1}}{2}, \text{ or } \frac{1}{2}$$

$$\sin 45^{\circ} = \frac{\sqrt{2}}{2}$$

$$\sin 60^{\circ} = \frac{\sqrt{3}}{2}$$

$$\sin 90^{\circ} = \frac{\sqrt{4}}{2}, \text{ or } 1$$

A similar pattern exists for the cosine function, except the values are given in reverse order.

Because you know the exact trigonometric values of 30° , 45° , and 60° angles, you can find the exact trigonometric values of *all* angles for which these angles are reference angles. The table lists these values for θ in both degrees and radians.

θ	$30^\circ \text{ or } \frac{\pi}{6}$	45° or $\frac{\pi}{4}$	60° or $\frac{\pi}{3}$
$\sin \theta$	$\frac{1}{2}$	$\frac{\sqrt{2}}{2}$	$\frac{\sqrt{3}}{2}$
$\cos \theta$	$\frac{\sqrt{3}}{2}$	$\frac{\sqrt{2}}{2}$	$\frac{1}{2}$
$\tan \theta$	$\frac{\sqrt{3}}{3}$	1	$\sqrt{3}$

Example 4 Use Reference Angles to Find Trigonometric Values

Find the exact value of each expression.

a. cos 120°

Because the terminal side of θ lies in Quadrant II, the reference angle θ' is $180^\circ - 120^\circ$ or 60° .

 $\cos 120^\circ = -\cos 60^\circ$ In Quadrant II, $\cos \theta$ is negative.

$$=-\frac{1}{2}$$
 cos 60° =

b. $\tan \frac{7\pi}{6}$

Because the terminal side of θ lies in Quadrant III, the reference angle θ' is $\frac{7\pi}{6} - \pi$ or $\frac{\pi}{6}$.

$$\tan \frac{7\pi}{6} = \tan \frac{\pi}{6}$$
 In Quadrant III, $\tan \theta$ is positive.
 $= \frac{\sqrt{3}}{3}$ $\tan \frac{\pi}{6} = \frac{\sqrt{3}}{3}$







c. $\csc \frac{15\pi}{4}$

4 -

A coterminal angle of θ is $\frac{15\pi}{4} - 2\pi$ or $\frac{7\pi}{4}$, which lies in Quadrant IV. So, the reference angle θ' is $2\pi - \frac{7\pi}{4}$ or $\frac{\pi}{4}$. Because sine and cosecant are reciprocal functions and sin θ is negative in Quadrant IV, it follows that $\csc \theta$ is also negative in Quadrant IV.



$$\csc \frac{15\pi}{4} = -\csc \frac{\pi}{4} \qquad \text{In Quadrant IV, } \csc \theta \text{ is negative}$$
$$= -\frac{1}{\sin \frac{\pi}{4}} \qquad \csc \theta = \frac{1}{\sin \theta}$$
$$= -\frac{1}{\frac{\sqrt{2}}{2}} \text{ or } -\sqrt{2} \qquad \sin \frac{\pi}{4} = \frac{\sqrt{2}}{2}$$

CHECK You can check your answer by using a graphing calculator.

$$\csc \frac{15\pi}{4} \approx -1.414 \checkmark$$
$$-\sqrt{2} \approx -1.414 \checkmark$$

GuidedPractice

Find the exact value of each expression.

4A. $\tan \frac{5\pi}{3}$ **4B.** $\sin \frac{5\pi}{6}$ **4C.** $\sec(-135^{\circ})$

If the value of one or more of the trigonometric functions and the quadrant in which the terminal side of θ lies is known, the remaining function values can be found.

Example 5 Use One Trigonometric Value to Find Others

Let $\tan \theta = \frac{5}{12}$, where $\sin \theta < 0$. Find the exact values of the five remaining trigonometric functions of θ .

To find the other function values, you must find the coordinates of a point on the terminal side of θ . You know that tan θ is positive and sin θ is negative, so θ must lie in Quadrant III. This means that both *x* and *y* are negative.

Because
$$\tan \theta = \frac{y}{x}$$
 or $\frac{5}{12}$, use the point (-12, -5) to find *r*.
 $r = \sqrt{x^2 + y^2}$ Pythagorean Theorem
 $= \sqrt{(-12)^2 + (-5)^2}$ $x = -12$ and $y = -5$
 $= \sqrt{169}$ or 13 Take the positive square root.

Use x = -12, y = -5, and r = 13 to write the five remaining trigonometric ratios.

$$\sin \theta = \frac{y}{r} \text{ or } -\frac{5}{13} \qquad \qquad \cos \theta = \frac{x}{r} \text{ or } -\frac{12}{13} \qquad \qquad \cot \theta = \frac{x}{y} \text{ or } \frac{12}{5}$$
$$\csc \theta = \frac{r}{y} \text{ or } -\frac{13}{5} \qquad \qquad \sec \theta = \frac{r}{x} \text{ or } -\frac{13}{12}$$

GuidedPractice

Find the exact values of the five remaining trigonometric functions of θ . **5A.** sec $\theta = \sqrt{3}$, tan $\theta < 0$ **5B.** sin $\theta = \frac{5}{7}$, cot $\theta > 0$

Rationalizing the Denominator Be sure to rationalize the denominator, if necessary.

Real-World Example 6 Find Coordinates Given a Radius and an Angle

ROBOTICS As part of the range of motion category in a high school robotics competition, a student programmed a 20-centimeter long robotic arm to pick up an object at point C and rotate through an angle of exactly 225° in order to release it into a container at point D. Find the position of the object at point D, relative to the pivot point O.



With the pivot point at the origin and the angle through which the arm rotates in standard position, point *C* has coordinates (20, 0). The reference angle θ' for 225° is 225° – 180° or 45°.

Let the position of point *D* have coordinates (x, y). The definitions of sine and cosine can then be used to find the values of *x* and *y*. The value of *r*, 20 centimeters, is the length of the robotic arm. Since *D* is in Quadrant III, the sine and cosine of 225° are negative.

$\cos\theta = \frac{x}{r}$	Cosine ratio	$\sin \theta = \frac{y}{r}$	Sine ratio
$\cos 225^\circ = \frac{x}{20}$	$\theta = 225^\circ$ and $r = 20$	$\sin 225^\circ = \frac{y}{20}$	$\theta = 225^\circ$ and $r = 20^\circ$
$-\cos 45^\circ = \frac{x}{20}$	$\cos 225^\circ = -\cos 45^\circ$	$-\sin 45^\circ = \frac{y}{20}$	$\sin 225^\circ = -\sin 45^\circ$
$-\frac{\sqrt{2}}{2} = \frac{x}{20}$	$\cos 45^\circ = \frac{\sqrt{2}}{2}$	$-\frac{\sqrt{2}}{2} = \frac{y}{20}$	$\sin 45^\circ = \frac{\sqrt{2}}{2}$
$-10\sqrt{2} = x$	Solve for <i>x</i> .	$-10\sqrt{2} = y$	Solve for y.

The exact coordinates of *D* are $(-10\sqrt{2}, -10\sqrt{2})$. Since $10\sqrt{2}$ is about 14.14, the object is about 14.14 centimeters to the left of the pivot point and about 14.14 centimeters below the pivot point.

GuidedPractice

6. CLOCKWORK A 3-inch-long minute hand on a clock shows a time of 45 minutes past the hour. What is the new position of the end of the minute hand relative to the pivot point at 10 minutes past the next hour?



Trigonometric Functions on the Unit Circle A

unit circle is a circle of radius 1 centered at the origin. Notice thaton a unit circle, the radian measure of a central angle $\theta = \frac{s}{1}$ or *s*, so the arc length intercepted by θ corresponds exactly to the angle's radian measure. This provides a way of mapping a real number input value for a trigonometric function to a real number output value.



StudyTip

Wrapping Function The

association of a point on the number line with a point on a circle is called the *wrapping function*, w(t). For example, if w(t) associates a point *t* on the number line with a point P(x, y) on the unit circle, then $w(\pi) = (-1, 0)$ and $w(2\pi) = (1, 0)$.

Consider the real number line placed vertically tangent to the unit circle at (1, 0) as shown below. If this line were wrapped about the circle in both the positive (counterclockwise) and negative (clockwise) direction, each point *t* on the line would map to a unique point *P*(*x*, *y*) on the circle. Because *r* = 1, we can define the trigonometric ratios of angle *t* in terms of just *x* and *y*.

Positive Values of t







Real-WorldLink

RoboCup is an international competition in which teams compete in a series of soccer matches, depending on the size and intelligence of their robots. The aim of the project is to advance artificial intelligence and robotics research.

Source: RoboCup

KeyConcept Trigonometric Functions on the Unit Circle

Let *t* be any real number on a number line and let P(x, y) be the point on *t* when the number line is wrapped onto the unit circle. Then the trigonometric functions of *t* are as follows. P(x, y) or $P(\cos t, \sin t)$



Notice that the input value in each of the definitions above can be thought of as an angle measure or as a real number *t*. When defined as functions of the real number system using the unit circle, the trigonometric functions are often called **circular functions**.

Using reference angles or quadrantal angles, you should now be able to find the trigonometric function values for all integer multiples of 30°, or $\frac{\pi}{6}$ radians, and 45°, or $\frac{\pi}{4}$ radians. These special values wrap to 16 special points on the unit circle, as shown below.



StudyTip

16-Point Unit Circle You have already memorized these values in the first quadrant. The remaining values can be determined using the *x*-axis, *y*-axis, and origin symmetry of the unit circle along with the signs of *x* and *y* in each quadrant.

Using the (x, y) coordinates in the 16-point unit circle and the definitions in the Key Concept Box at the top of the page, you can find the values of the trigonometric functions for common angle measures. It is helpful to memorize these exact function values so you can quickly perform calculations involving them.

Example 7 Find Trignometric Values Using the Unit Circle

Find the exact value of each expression. If undefined, write undefined.

a. $\sin \frac{\pi}{3}$ $\frac{\pi}{3}$ corresponds to the point $(x, y) = \left(\frac{1}{2}, \frac{\sqrt{3}}{2}\right)$ on the unit circle. $\sin t = y$ Definition of $\sin t$ $\sin \frac{\pi}{3} = \frac{\sqrt{3}}{2}$ $y = \frac{\sqrt{3}}{2}$ when $t = \frac{\pi}{3}$.

b. cos 135°

135° corresponds to the point $(x, y) = \left(-\frac{\sqrt{2}}{2}, \frac{\sqrt{2}}{2}\right)$ on the unit circle. $\cos t = x$ Definition of $\cos t$

 $\cos 135^\circ = -\frac{\sqrt{2}}{2}$ $x = -\frac{\sqrt{2}}{2}$ when $t = 135^\circ$.

c. tan 270°

270° corresponds to the point (x, y) = (0, -1) on the unit circle.

$$\tan t = \frac{y}{x}$$
 Definition of $\tan t$
$$\tan 270^\circ = \frac{-1}{0}$$
 $x = 0$ and $y = -1$, when $t = 270^\circ$.

Therefore, tan 270° is undefined.

d. $\csc \frac{11\pi}{6}$ $\frac{11\pi}{6}$ corresponds to the point $(x, y) = \left(\frac{\sqrt{3}}{2}, -\frac{1}{2}\right)$ on the unit circle. $\csc t = \frac{1}{y}$ Definition of $\csc t$ $\csc \frac{11\pi}{6} = \frac{1}{-\frac{1}{2}}$ $y = -\frac{1}{2}$ when $t = \frac{11\pi}{6}$. = -2 Simplify. GuidedPractice 7A. $\cos \frac{\pi}{4}$ 7B. $\sin 120^{\circ}$ 7C. $\cot 210^{\circ}$ 7D. $\sec \frac{7\pi}{4}$

StudyTip

Radians vs. Degrees While we could also discuss one wrapping as corresponding to an angle measure of 360° , this measure is not related to a distance. On the unit circle, one wrapping corresponds to both the angle measuring 2π and the distance 2π around the circle. As defined by wrapping the number line around the unit circle, the domain of both the sine and cosine functions is the set of all real numbers $(-\infty, \infty)$. Extending infinitely in either direction, the number line can be wrapped multiple times around the unit circle, mapping more than one *t*-value to the same point P(x, y) with each wrapping, positive or negative.



Because $\cos t = x$, $\sin t = y$, and one wrapping corresponds to a distance of 2π ,

 $\cos(t + 2n\pi) = \cos t$ and $\sin(t + 2n\pi) = \sin t$,

for any integer *n* and real number *t*.

StudyTip

Periodic Functions The other three circular functions are also periodic. The periods of these functions will be discussed in Lesson 4-5. The values for the sine and cosine function therefore lie in the interval [-1, 1] and repeat for every integer multiple of 2π on the number line. Functions with values that repeat at regular intervals are called **periodic functions**.

KeyConcept Periodic Functions

```
A function y = f(t) is periodic if there exists a positive real number c such that f(t + c) = f(t) for all values of t in the domain of f.
```

The smallest number *c* for which *f* is periodic is called the **period** of *f*.

The sine and cosine functions are periodic, repeating values after 2π , so these functions have a period of 2π . It can be shown that the values of the tangent function repeat after a distance of π on the number line, so the tangent function has a period of π and

$$\tan t = \tan \left(t + n\pi\right),$$

for any integer *n* and real number *t*, unless both tan *t* and tan $(t + n\pi)$ are undefined. You can use the periodic nature of the sine, cosine, and tangent functions to evaluate these functions.

Example 8 Use the Periodic Nature of Circular Functions Find the exact value of each expression. a. $\cos \frac{11\pi}{4}$ $\cos \frac{11\pi}{4} = \cos \left(\frac{3\pi}{4} + 2\pi \right)$ Rewrite $\frac{11\pi}{4}$ as the sum of a number and 2π . $\frac{3\pi}{4}$ and $\frac{3\pi}{4} + 2\pi$ map to the same point $(x, y) = \left(-\frac{\sqrt{2}}{2}, \frac{\sqrt{2}}{2}\right)$ on the unit circle. $=\cos\frac{3\pi}{4}$ $=-\frac{\sqrt{2}}{2}$ $\cos t = x$ and $x = -\frac{\sqrt{2}}{2}$ when $t = \frac{3\pi}{4}$. b. sin $\left(-\frac{2\pi}{2}\right)$ $\sin\left(-\frac{2\pi}{3}\right) = \sin\left(\frac{4\pi}{3} + 2(-1)\pi\right)$ Rewrite $-\frac{2\pi}{3}$ as the sum of a number and an integer multiple of 2π . $\frac{4\pi}{3}$ and $\frac{4\pi}{3} - 2(-1)\pi$ map to the same point (x, y) = $\left(-\frac{1}{2}, -\frac{\sqrt{3}}{2}\right)$ on $=\sin\frac{4\pi}{2}$ $=-\frac{\sqrt{3}}{2}$ sin t = y and $y = -\frac{\sqrt{3}}{2}$ when $t = \frac{4\pi}{3}$ **c.** $\tan \frac{19\pi}{6}$ $\tan \frac{19\pi}{6} = \tan \left(\frac{\pi}{6} + 3\pi\right)$ Rewrite $\frac{19\pi}{6}$ as the sum of a number and an integer multiple of π . $= \tan \frac{\pi}{6}$ $\frac{\pi}{6}$ and $\frac{\pi}{6}+3\pi$ map to points on the unit circle with the same $=\frac{\frac{1}{2}}{\frac{\sqrt{3}}{3}} \text{ or } \frac{\sqrt{3}}{3}$ tan $t = \frac{y}{x}$; $x = \frac{\sqrt{3}}{2}$ and $y = \frac{1}{2}$ when $t = \frac{\pi}{6}$. GuidedPractice **8B.** $\cos\left(-\frac{4\pi}{2}\right)$ **8A.** $\sin \frac{13\pi}{4}$ **8C.** $\tan \frac{15\pi}{6}$

Recall from Lesson 1-2 that a function *f* is *even* if for every *x* in the domain of *f*, f(-x) = f(x) and *odd* if for every *x* in the domain of *f*, f(-x) = -f(x). You can use the unit circle to verify that the cosine function is even and that the sine and tangent functions are odd. That is,

 $\cos(-t) = \cos t$ $\sin(-t) = -\sin t$ $\tan(-t) = -\tan t$.

Exercises

The given point lies on the terminal side of an angle θ in standard position. Find the values of the six trigonometric functions of θ . (Example 1)

1.	(3, 4)	2.	(-6,6)
3.	(-4, -3)	4.	(2, 0)
5.	(1, -8)	6.	(5, -3)
7.	(-8, 15)	8.	(-1, -2)

Find the exact value of each trigonometric function, if defined. If not defined, write *undefined*. (Example 2)

9.	$\sin\frac{\pi}{2}$	10.	tan 2 π
11.	cot (-180°)	12.	$\csc 270^{\circ}$
13.	cos (-270°)	14.	sec 180°
15.	$\tan \pi$	16.	$\sec\left(-\frac{\pi}{2}\right)$

Sketch each angle. Then find its reference angle. (Example 3)

17.	135°	18.	210°
19.	$\frac{7\pi}{12}$	20.	$\frac{11\pi}{3}$
21.	-405°	22.	-75°
23.	$\frac{5\pi}{6}$	24.	$\frac{13\pi}{6}$

Find the exact value of each expression. (Example 4)

25.	$\cos\frac{4\pi}{3}$	26.	$\tan \frac{7\pi}{6}$
27.	$\sin \frac{3\pi}{4}$	28.	cot (-45°)
29.	csc 390°	30.	sec (-150°)
31.	$\tan \frac{11\pi}{6}$	32.	sin 300°

Find the exact values of the five remaining trigonometric functions of θ . (Example 5)

- **33.** tan $\theta = 2$, where sin $\theta > 0$ and cos $\theta > 0$
- **34.** csc θ = 2, where sin θ > 0 and cos θ < 0
- **35.** $\sin \theta = -\frac{1}{5}$, where $\cos \theta > 0$
- **36.** $\cos \theta = -\frac{12}{13}$, where $\sin \theta < 0$
- **37.** sec $\theta = \sqrt{3}$, where sin $\theta < 0$ and cos $\theta > 0$
- **38.** $\cot \theta = 1$, where $\sin \theta < 0$ and $\cos \theta < 0$
- **39.** $\tan \theta = -1$, where $\sin \theta < 0$

40.
$$\cos \theta = -\frac{1}{2}$$
, where $\sin \theta > 0$

41 CAROUSEL Zoe is on a carousel at the carnival. The diameter of the carousel is 80 feet. Find the position of her seat from the center of the carousel after a rotation of 210°. (Example 6)



42. COIN FUNNEL A coin is dropped into a funnel where it spins in smaller circles until it drops into the bottom of the bank. The diameter of the first circle the coin makes is 24 centimeters. Before completing one full circle, the coin travels 150° and falls over. What is the new position of the coin relative to the center of the funnel? (Example 6)

Find the exact value of each expression. If undefined, write *undefined*. (Examples 7 and 8)

43.	sec 120°	44.	sin 315°
45.	$\cos\frac{11\pi}{3}$	46.	$\tan\left(-\frac{5\pi}{4}\right)$
47.	csc 390°	48.	$\cot 510^{\circ}$
49.	csc 5400°	50.	$\sec \frac{3\pi}{2}$
51.	$\cot\left(-\frac{5\pi}{6}\right)$	52.	$\csc \frac{17\pi}{6}$
53.	$\tan\frac{5\pi}{3}$	54.	$\sec \frac{7\pi}{6}$
55.	$\sin\left(-\frac{5\pi}{3}\right)$	56.	$\cos\frac{7\pi}{4}$
57.	$\tan\frac{14\pi}{3}$	58.	$\cos\left(-\frac{19\pi}{6}\right)$

59. RIDES Jae and Anya are on a ride at an amusement park. After the first several swings, the angle the ride makes with the vertical is modeled by $\theta = 22 \cos \pi t$, with θ measured in radians and *t* measured in seconds. Determine the measure of the angle in radians for t = 0, 0.5, 1, 1.5, 2, and 2.5. (Example 8)





Complete each trigonometric expression.

60.	$\cos 60^\circ = \sin $	61.	$\tan\frac{\pi}{4} = \sin$
62.	$\sin\frac{2\pi}{3} = \cos$	63.	$\cos\frac{7\pi}{6} = \sin$
64.	$\sin\left(-45^\circ\right) = \cos$	65.	$\cos\frac{5\pi}{3} = \sin$

- **66. ICE CREAM** The monthly sales in thousands of dollars for Fiona's Fine Ice Cream shop can be modeled by $y = 71.3 + 59.6 \sin \frac{\pi(t-4)}{6}$, where t = 1 represents January, t = 2 represents February, and so on.
 - **a.** Estimate the sales for January, March, July, and October.
 - **b.** Describe why the ice cream shop's sales can be represented by a trigonometric function.

Use the given values to evaluate the trigonometric functions.

- **67.** $\cos(-\theta) = \frac{8}{11}; \cos \theta = ?; \sec \theta = ?$
- **68.** $\sin(-\theta) = \frac{5}{9}; \sin \theta = ?; \csc \theta = ?$
- **69.** $\sec \theta = \frac{13}{12}; \cos \theta = ?; \cos(-\theta) = ?$
- **70.** $\csc \theta = \frac{19}{17}; \sin \theta = ?; \sin(-\theta) = ?$
- **71. GRAPHS** Suppose the terminal side of an angle θ in standard position coincides with the graph of y = 2x in Quadrant III. Find the six trigonometric functions of θ .

Find the coordinates of *P* for each circle with the given radius and angle measure.



76. COMPARISON Suppose the terminal side of an angle θ_1 in standard position contains the point (7, -8), and the terminal side of a second angle θ_2 in standard position contains the point (-7, 8). Compare the sines of θ_1 and θ_2 .

77. TIDES The depth *y* in meters of the tide on a beach varies as a sine function of *x*, the hour of the day. On a certain

day, that function was $y = 3 \sin \left[\frac{\pi}{6}(x-4)\right] + 8$, where

x = 0, 1, 2, ..., 24 corresponds to 12:00 midnight, 1:00 а.м., 2:00 а.м., ..., 12:00 midnight the next night.

- **a.** What is the maximum depth, or high tide, that day?
- **b.** At what time(s) does the high tide occur?
- **78. MULTIPLE REPRESENTATIONS** In this problem, you will investigate the period of the sine function.
 - **a. TABULAR** Copy and complete a table similar to the one below that includes all 16 angle measures from the unit circle.

θ	0	$\frac{\pi}{6}$	$\frac{\pi}{4}$	$\frac{\pi}{3}$	 2π
sin <i>θ</i>					
sin 2 <i>0</i>					
sin 4 <i>0</i>					

- **b. VERBAL** After what values of *θ* do sin *θ*, sin 2*θ*, and sin 4*θ*, repeat their range values? In other words, what are the periods of these functions?
- **c. VERBAL** Make a conjecture as to how the period of $y = \sin n\theta$ is affected for different values of *n*.

H.O.T. Problems Use Higher-Order Thinking Skills

79 CHALLENGE For each statement, describe *n*.

- **a.** $\cos\left(n\cdot\frac{\pi}{2}\right)=0$
- **b.** $\csc\left(n \cdot \frac{\pi}{2}\right)$ is undefined.

REASONING Determine whether each statement is *true* or *false*. Explain your reasoning.

- **80.** If $\cos \theta = 0.8$, $\sec \theta \cos (-\theta) = 0.45$.
- **81.** Since $\tan(-t) = -\tan t$, the tangent of a negative angle is a negative number.
- **82.** WRITING IN MATH Explain why the attendance at a year-round theme park could be modeled by a periodic function. What issues or events could occur over time to alter this periodic depiction?

REASONING Use the unit circle to verify each relationship.

- **83.** $\sin(-t) = -\sin t$
- **84.** $\cos(-t) = \cos t$
- **85.** $\tan(-t) = -\tan t$
- **86.** WRITING IN MATH Make a conjecture as to the periods of the secant, cosecant, and cotangent functions. Explain your reasoning.

Spiral Review

Write each decimal degree measure in DMS form and each DMS measure in decimal degree form to the nearest thousandth. (Lesson 4-2)

87. 168.35° 88. 27.465°	89. 14° 5′ 20″	90. 173° 1
---------------------------------------	-----------------------	-------------------

24'35"



b. If the treadmill bed is 40 inches long, what is the vertical rise when set at an 8% incline?

when set at a 10% incline? Round to the nearest degree.

Evaluate each logarithm. (Lesson 3-3)

for every 100 units of horizontal run. (Lesson 4-1)

92. $\log_8 64$ 93. $\log_{125} 5$ 94. $\log_2 32$ 95. $\log_4 1$	92. log ₈ 64	93. log ₁₂₅ 5	94. log ₂ 32	95. log ₄ 128
---	--------------------------------	---------------------------------	--------------------------------	---------------------------------

List all possible rational zeros of each function. Then determine which, if any, are zeros. (Lesson 2-4)

96. $f(x) = x^3 - 4x^2 + x + 2$	97. $g(x) = x^3 + 6x^2 + 10x + 3$
98. $h(x) = x^4 - x^2 + x - 1$	99. $h(x) = 2x^3 + 3x^2 - 8x + 3$
100. $f(x) = 2x^4 + 3x^3 - 6x^2 - 11x - 3$	101. $g(x) = 4x^3 + x^2 + 8x + 2$

102. NAVIGATION A global positioning system (GPS) uses satellites to allow a user to determine his or her position on Earth. The system depends on satellite signals that are reflected to and from a hand-held transmitter. The time that the signal takes to reflect is used to determine the transmitter's position. Radio waves travel through air at a speed of 299,792,458 meters per second. Thus, d(t) = 299,792,458t relates the time t in seconds to the distance traveled d(t) in meters. (Lesson 1-1)

- **a.** Find the distance a radio wave will travel in 0.05, 0.2, 1.4, and 5.9 seconds.
- **b.** If a signal from a GPS satellite is received at a transmitter in 0.08 second, how far from the transmitter is the satellite?

Skills Review for Standardized Tests

103. SAT/ACT In the figure, \overline{AB} and \overline{AD} are tangents to circle *C*. What is the value of *m*?



- **104.** Suppose θ is an angle in standard position with $\sin \theta > 0$. In which quadrant(s) could the terminal side of θ lie?
 - C I and III A I only D I and IV **B** I and II

- 105. **REVIEW** Find the angular speed in radians per second of a point on a bicycle tire if it completes 2 revolutions in 3 seconds.
 - F $\frac{\pi}{3}$
 - G
 - H $\frac{2\pi}{3}$
 - J $\frac{4\pi}{3}$
- **106. REVIEW** Which angle has a tangent and cosine that are both negative?
 - A 110°
 - **B** 180°
 - C 210°
 - **D** 340°


Graphing Technology Lab Graphing the Sine Function Parametrically



Objective

 Use a graphing calculator and parametric equations to graph the sine function and its inverse. As functions of the real number system, you can graph trigonometric functions on the coordinate plane and apply the same graphical analysis that you did to functions in Chapter 1. As was done in Extend 1-7, parametric equations will be used to graph the sine function.

Activity 1 Parametric Graph of $y = \sin x$





Step 1 Set the mode. In the MODE menu, select RADIAN, PAR, and SIMUL. This allows the equations to be graphed simultaneously. Next, enter the parametric equations. In parametric form, X,T,θ,n will use *t* instead of *x*.

TURIAL SCI ENG
RHDIAN DEGREE
FUNC PAR POL SEQ
CONGECTED DOT
SEQUEDTIAL SECUE
arbi refei
FULL BORIZ G-T
SET CLOCK



Step 2 Set the *x*- and *t*-values to range from 0 to 2π . Set Tstep and *x*-scale to $\frac{\pi}{12}$. Set *y* to [-1, 1] scl: 0.1. The calculator automatically converts to decimal form.

Step 3 Graph the equations. Trace the function to identify points along the graph. Select **TRACE** and use the right arrow to move along the curve. Record the corresponding *x*- and *y*-values.





 $t: [0, 2\pi]; t \text{ step } \frac{\pi}{12}$

Step 4 The table shows angle measures from 0° to 180° , or 0 to π , and the corresponding values for sin *t* on the unit circle. The figures below illustrate the relationship between the graph and the unit circle.

Degrees	0	30	45	60	90	120	135	150	180
Radians	0	0.52	0.79	1.05	1.571	2.094	2.356	2.618	3.14
$y = \sin t$	0	0.5	0.707	0.866	1	0.866	0.707	0.5	0





StudyTip

Decimal Equivalents Below are the decimal equivalents of common trigonometric values.



Exercises

Graph each function on $[0, 2\pi]$.

1. $x = t, y$	$=\cos t$	2.	$x = t, y = \sin 2t$
3. $x = t, y$	$= 3 \cos t$	4.	$x = t, y = 4\sin t$
5. $x = t, y$	$=\cos\left(t+\pi\right)$	6.	$x = t, y = 2\sin\left(t - \frac{\pi}{4}\right)$

By definition, sin t is the y-coordinate of the point P(x, y) on the unit circle to which the real number t on the number line gets wrapped. As shown in the diagram on the previous page, the graph of $y = \sin t$ follows the y-coordinate of the point determined by t as it moves counterclockwise around the unit circle.

The graph of the sine function is called a sine curve. From Lesson 4-3, you know that the sine function is periodic with a period of 2π . That is, the sine curve graphed from 0 to 2π would repeat every distance of 2π in either direction, positive or negative. Parametric equations can be used to graph the inverse of the sine function.



Exercises

StudyTip

smoother curve.

Graph each function and its inverse. Then determine a domain for which each function is one-to-one.

7. $x = t, y = \cos 2t$	8. $x = t, y = -\sin t$
9. $x = t, y = 2 \cos t$	10. $x = t + \frac{\pi}{4}, y = \sin t$
11. $x = t, y = 2\cos(t - \pi)$	12. $x = t - \frac{\pi}{6}, y = \sin t$

Graphing Sine and Cosine Functions

Then

Now

Why?

- You analyzed graphs of functions.
- Graph transformations As you ride a Ferris wheel, the height that you are above the ground varies periodically as a function of time. You can model this behavior using a sinusoidal function.
- Use sinusoidal functions to solve problems.

functions.

of the sine and cosine



NewVocabulary

sinusoid amplitude frequency phase shift vertical shift midline

Transformations of Sine and Cosine Functions As shown in Explore 4-4, the graph $y = \sin t$ follows the y-coordinate of the point determined by t as it moves around the unit circle. Similarly, the graph of $y = \cos t$ follows the *x*-coordinate of this point. The graphs of these functions are periodic, repeating after a period of 2π . The properties of the sine and cosine functions are summarized below.



The portion of each graph on $[0, 2\pi]$ represents one period or *cycle* of the function. Notice that the cosine graph is a horizontal translation of the sine graph. Any transformation of a sine function is called a sinusoid. The general form of the sinusoidal functions sine and cosine are

$$y = a \sin(bx + c) + d$$
 and $y = a \cos(bx + c) + d$

where *a*, *b*, *c*, and *d* are constants and neither *a* nor *b* is 0.

Notice that the constant factor a in $y = a \sin x$ and $y = a \cos x$ expands the graphs of $y = \sin x$ and $y = \cos x$ vertically if |a| > 1 and compresses them vertically if |a| < 1.

StudyTip

Dilations and x-intercepts Notice that a dilation of a sinusoidal function does not affect where the curve crosses the x-axis, at its x-intercepts.



Vertical dilations affect the amplitude of sinusoidal functions.



To graph a sinusoidal function of the form $y = a \sin x$ or $y = a \cos x$, plot the x-intercepts of the parent sine or cosine function and use the amplitude |a| to plot the new maximum and minimum points. Then sketch the sine wave through these points.

Example 1 Graph Vertical Dilations of Sinusoidal Functions

Describe how the graphs of $f(x) = \sin x$ and $g(x) = \frac{1}{4} \sin x$ are related. Then find the amplitude of g(x), and sketch two periods of both functions on the same coordinate axes.

The graph of g(x) is the graph of f(x) compressed vertically. The amplitude of g(x) is $\left|\frac{1}{4}\right|$ or $\frac{1}{4}$.

Create a table listing the coordinates of the *x*-intercepts and extrema for $f(x) = \sin x$ for one period on $[0, 2\pi]$. Then use the amplitude of g(x) to find corresponding points on its graph.

Function	<i>x</i> -intercept	Maximum	<i>x</i> -intercept	Minimum	<i>x</i> -intercept	
$f(x) = \sin x$	(0, 0)	$\left(\frac{\pi}{2},1\right)$	(π, 0)	$\left(\frac{3\pi}{2}, -1\right)$	(2π, 0)	
$g(x) = \frac{1}{4}\sin x$	(0, 0)	$\left(\frac{\pi}{2},\frac{1}{4}\right)$	(π, 0)	$\left(\frac{3\pi}{2},-\frac{1}{4}\right)$	(2π, 0)	

Sketch the curve through the indicated points for each function. Then repeat the pattern suggested by one period of each graph to complete a second period on $[2\pi, 4\pi]$. Extend each curve to the left and right to indicate that the curve continues in both directions.



GuidedPractice

Describe how the graphs of f(x) and g(x) are related. Then find the amplitude of g(x), and sketch two periods of both functions on the same coordinate axes.

1A.
$$f(x) = \cos x$$

 $g(x) = \frac{1}{3}\cos x$
1B. $f(x) = \sin x$
1C. $f(x) = \cos x$
 $g(x) = 2\cos x$

StudyTip

Radians Versus Degrees

You could rescale the x-axis in terms of degrees and produce sinusoidal graphs that look similar to those produced using radian measure. In calculus, however, you will encounter rules that depend on radian measure. So, in this book, we will graph all trigonometric functions in terms of radians.

x



Example 2 Graph Reflections of Sinusoidal Functions

Describe how the graphs of $f(x) = \cos x$ and $g(x) = -3 \cos x$ are related. Then find the amplitude of g(x), and sketch two periods of both functions on the same coordinate axes.

The graph of g(x) is the graph of f(x) expanded vertically and then reflected in the *x*-axis. The amplitude of g(x) is |-3| or 3.

Create a table listing the coordinates of key points of $f(x) = \cos x$ for one period on $[0, 2\pi]$. Use the amplitude of g(x) to find corresponding points on the graph of $y = 3 \cos x$. Then reflect these points in the *x*-axis to find corresponding points on the graph of g(x).

Function	Extremum	<i>x</i> -intercept	Extremum	<i>x</i> -intercept	Extremum
$f(x) = \cos x$	(0, 1)	$\left(\frac{\pi}{2},0\right)$	(π, -1)	$\left(\frac{3\pi}{2},0\right)$	(2π, 1)
$y = 3 \cos x$	(0, 3)	$\left(\frac{\pi}{2},0\right)$	(π, -3)	$\left(\frac{3\pi}{2},0\right)$	(2π, 3)
$g(x) = -3\cos x$	(0, —3)	$\left(\frac{\pi}{2},0\right)$	(π, 3)	$\left(\frac{3\pi}{2},0\right)$	(2π, -3)

Sketch the curve through the indicated points for each function. Then repeat the pattern suggested by one period of each graph to complete a second period on $[2\pi, 4\pi]$. Extend each curve to the left and right to indicate that the curve continues in both directions.



GuidedPractice

Describe how the graphs of f(x) and g(x) are related. Then find the amplitude of g(x), and sketch two periods of both functions on the same coordinate axes.

2A.
$$f(x) = \cos x$$

 $g(x) = -\frac{1}{5}\cos x$
2B. $f(x) = \sin x$
 $g(x) = -4\sin x$

In Lesson 1-5, you learned that if g(x) = f(bx), then g(x) is the graph of f(x) compressed horizontally if |b| > 1 and expanded horizontally if |b| < 1. Horizontal dilations affect the *period* of a sinusoidal function—the length of one full cycle.



WatchOut!

Amplitude Notice that Example 2 does not state that the amplitude of $g(x) = -3 \cos x$ is -3. Amplitude is a height and is not directional.



To graph a sinusoidal function of the form $y = \sin bx$ or $y = \cos bx$, find the period of the function and successively add $\frac{\text{period}}{4}$ to the left endpoint of an interval with that length. Then use these values as the *x*-values for the key points on the graph.

Example 3 Graph Horizontal Dilations of Sinusoidal Functions

Describe how the graphs of $f(x) = \cos x$ and $g(x) = \cos \frac{x}{3}$ are related. Then find the period of g(x), and sketch at least one period of both functions on the same coordinate axes.

Because $\cos \frac{x}{3} = \cos \frac{1}{3}x$, the graph of g(x) is the graph of f(x) expanded horizontally. The period of g(x) is $\frac{2\pi}{\left|\frac{1}{3}\right|}$ or 6π .

Because the period of g(x) is 6π , to find corresponding points on the graph of g(x), change the *x*-coordinates of those key points on f(x) so that they range from 0 to 6π , increasing by increments of $\frac{6\pi}{4}$ or $\frac{3\pi}{2}$.

F	unction	Maximum	<i>x</i> -intercept	Minimum	<i>x</i> -intercept	Maximum	
<i>f</i> (<i>x</i>)	$) = \cos x$	(<mark>0</mark> , 1)	$\left(\frac{\pi}{2},0\right)$	(π , −1)	$\left(\frac{3\pi}{2},0\right)$	(2 π, 1)	
g(x)	$r = \cos \frac{x}{3}$	(0 , 1)	$\left(\frac{3\pi}{2},0\right)$	(3 π, −1)	$\left(\frac{9\pi}{2},0\right)$	(6 π, 1)	

Sketch the curve through the indicated points for each function, continuing the patterns to complete one full cycle of each.



GuidedPractice

Describe how the graphs of f(x) and g(x) are related. Then find the period of g(x), and sketch at least one period of each function on the same coordinate axes.

3A. $f(x) = \cos x$	3B. $f(x) = \sin x$	3C. $f(x) = \cos x$
$g(x) = \cos\frac{x}{2}$	$g(x) = \sin 3x$	$g(x) = \cos\frac{1}{4}x$



Because the frequency of a sinusoidal function is the reciprocal of the period, it follows that the **p**eriod of the function is the reciprocal of its frequency.

Real-World Example 4 Use Frequency to Write a Sinusoidal Function

MUSIC Musical notes are classified by frequency. In the equal tempered scale, middle C has a frequency of 262 hertz. Use this information and the information at the left to write an equation for a sine function that can be used to model the initial behavior of the sound wave associated with middle C having an amplitude of 0.2.

The general form of the equation will be $y = a \sin bt$, where *t* is the time in seconds. Because the amplitude is 0.2, |a| = 0.2. This means that $a = \pm 0.2$.

The period is the reciprocal of the frequency or $\frac{1}{262}$. Use this value to find *b*.

period = $\frac{2\pi}{ b }$	Period formula
$\frac{1}{262} = \frac{2\pi}{ b }$	period = $\frac{1}{262}$
$ b = 2\pi(262)$ or 524π	Solve for b .
$b = \pm 524\pi$	Solve for <i>b</i> .

By arbitrarily choosing the positive values of *a* and *b*, one sine function that models the initial behavior is $y = 0.2 \sin 524\pi t$.

GuidedPractice

I

4. MUSIC In the same scale, the C above middle C has a frequency of 524 hertz. Write an equation for a sine function that can be used to model the initial behavior of the sound wave associated with this C having an amplitude of 0.1.

A *phase* of a sinusoid is the position of a wave relative to some reference point. A horizontal translation of a sinusoidal function results in a *phase shift*. Recall from Lesson 1-5 that the graph of y = f(x + c) is the graph of y = f(x) translated or *shifted* |c| units left if c > 0 and |c| units right if c < 0.



Real-WorldLink

In physics, frequency is measured

in *hertz* or oscillations per second. For example, the number of sound

waves passing a point A in one

second would be the wave's

Source: Science World

frequency.





StudyTip

Alternative Form The general forms of the sinusoidal functions can also be expressed as $y = a \sin b(x - h) + k$ and $y = a \cos b(x - h) + k$. In these forms, each sinusoid has a phase shift of *h* and a vertical translation of *k* in comparison to the graphs of $y = a \sin bx$ and $y = a \cos bx$. To graph the phase shift of a sinusoidal function of the form $y = a \sin(bx + c) + d$ or $y = a \cos(bx + c) + d$, first determine the endpoints of an interval that corresponds to one cycle of the graph by adding $-\frac{c}{h}$ to each endpoint on the interval $[0, 2\pi]$ of the parent function.

Example 5 Graph Horizontal Translations of Sinusoidal Functions

State the amplitude, period, frequency, and phase shift of $y = \sin\left(3x - \frac{\pi}{2}\right)$. Then graph two periods of the function.

In this function, a = 1, b = 3, and $c = -\frac{\pi}{2}$.

Amplitude: $|\mathbf{a}| = |\mathbf{1}|$ or 1

Frequency: $\frac{|b|}{2\pi} = \frac{|3|}{2\pi} \text{ or } \frac{3}{2\pi}$

Period:
$$\frac{2\pi}{|b|} = \frac{2\pi}{|3|}$$
 or $\frac{2\pi}{3}$
Phase shift: $-\frac{c}{|b|} = -\frac{-\frac{\pi}{2}}{|3|}$ or $\frac{\pi}{6}$

To graph $y = \sin\left(3x - \frac{\pi}{2}\right)$, consider the graph of $y = \sin 3x$. The period of this function is $\frac{2\pi}{3}$. Create a table listing the coordinates of key points of $y = \sin 3x$ on the interval $\left[0, \frac{2\pi}{3}\right]$. To account for a phase shift of $\frac{\pi}{6}$, add $\frac{\pi}{6}$ to the *x*-values of each of the key points for the graph of $y = \sin 3x$.

Function	<i>x</i> -intercept	Maximum	<i>x</i> -intercept	Minimum	<i>x-</i> intercept	
$y = \sin 3x$	(0 , 0)	$\left(\frac{\pi}{6},1\right)$	$\left(\frac{\pi}{3},0\right)$	$\left(\frac{\pi}{2},-1\right)$	$\left(\frac{2\pi}{3},0\right)$	
$y = \sin\left(3x - \frac{\pi}{2}\right)$	$\left(\frac{\pi}{6},0\right)$	$\left(\frac{\pi}{3},1\right)$	$\left(\frac{\pi}{2},0\right)$	$\left(\frac{2\pi}{3},-1\right)$	$\left(\frac{5\pi}{6}, 0\right)$	

Sketch the graph of $y = \sin\left(3x - \frac{\pi}{2}\right)$ through these points, continuing the pattern to complete two cycles.



GuidedPractice

State the amplitude, period, frequency, and phase shift of each function. Then graph two periods of the function.

5A.
$$y = \cos\left(\frac{x}{2} + \frac{\pi}{4}\right)$$

5B.
$$y = 3 \sin \left(2x - \frac{\pi}{3}\right)$$

StudyTip

Notation $\sin(x+d) \neq \sin x + d$ The first expression indicates a phase shift, while the second expression indicates a vertical shift.

The final way to transform the graph of a sinusoidal function is through a vertical translation or **vertical shift**. Recall from Lesson 1-5 that the graph of y = f(x) + d is the graph of y = f(x)translated or *shifted* |d| units up if d > 0 and |d| units down if d < 0. The vertical shift is the average of the maximum and minimum values of the function.

The parent functions $y = \sin x$ and $y = \cos x$ oscillate about the *x*-axis. After a vertical shift, a new horizontal axis known as the **midline** becomes the reference line or equilibrium point about which the graph oscillates. For example, the midline of $y = \sin x + 1$ is y = 1, as shown.



In general, the midline for the graphs of $y = a \sin(bx + c) + d$ and $y = a \cos(bx + c) + d$ is y = d.

Example 6 Graph Vertical Translations of Sinusoidal Functions

State the amplitude, period, frequency, phase shift, and vertical shift of $y = \sin(x + 2\pi) - 1$. Then graph two periods of the function.

In this function, a = 1, b = 1, $c = 2\pi$, and d = -1.

Amplitude: |a| = |1| or 1Period: $\frac{2\pi}{|b|} = \frac{2\pi}{|1|}$ or 2π Frequency: $\frac{|b|}{2\pi} = \frac{|1|}{2\pi}$ or $\frac{1}{2\pi}$ Phase shift: $-\frac{c}{|b|} = -\frac{2\pi}{|1|} = -2\pi$ Vertical shift: d or -1Midline: y = d or y = -1

First, graph the midline y = -1. Then graph $y = \sin x$ shifted 2π units to the left and 1 unit down.

Notice that this transformation is equivalent to a translation 1 unit down because the phase shift was one period to the left.



GuidedPractice

State the amplitude, period, frequency, phase shift, and vertical shift of each function. Then graph two periods of the function.

6A.
$$y = 2\cos x + 1$$

6B.
$$y = \frac{1}{2} \sin\left(\frac{x}{4} - \frac{\pi}{2}\right) - 3$$

The characteristics of transformations of the parent functions $y = \sin x$ and $y = \cos x$ are summarized below.

ConceptSummary Graphs of Sinusoidal Functions The graphs of $y = a \sin(bx + c) + d$ and $y = a \cos(bx + c) + d$, where $a \neq 0$ and $b \neq 0$, have the following characteristics. Frequency: $\frac{|b|}{2\pi}$ or $\frac{1}{\text{Period}}$ Amplitude: |a| Period: $\frac{2\pi}{|b|}$ Phase shift: $-\frac{c}{|b|}$ Vertical shift: d Midline: v = d

TechnologyTip

Zoom Trig When graphing a trigonometric function using your graphing calculator, be sure you are in radian mode and use the ZTrig selection under the zoom feature to change your viewing window from the standard window to a more appropriate window of $[-2\pi, 2\pi]$ scl: $\pi/2$ by [-4, 4] scl: 1.



Real-WorldLink

The table shows the number of daylight hours on the 15th of each month in New York City.

Month	Hours of Daylight
January	9.58
February	10.67
March	11.9
April	13.3
May	14.43
June	15.07
July	14.8
August	13.8
September	12.48
October	11.15
November	9.9
December	9.27
Source: U.S. Nav	al Observatory

[0, 12] scl: 1 by [0, 20] scl: 2

Figure 4.4.1

Real-World Example 7 Modeling Data Using a Sinusoidal Function

behavior over time can be modeled by transformations of $y = \sin x$ or $y = \cos x$.

METEOROLOGY Use the information at the left to write a sinusoidal function that models the number of hours of daylight for New York City as a function of time x, where x = 1 represents January 15, x = 2 represents February 15, and so on. Then use your model to estimate the number of hours of daylight on September 30 in New York City.

Applications of Sinusoidal Functions Many real-world situations that exhibit periodic

Step 1 Make a scatter plot of the data and choose a model.

The graph appears wave-like, so you can use a sinusoidal function of the form $y = a \sin(bx + c) + d$ or $y = a \cos(bx + c) + d$ to model the data. We will choose to use $y = a \cos(bx + c) + d$ to model the data.



Step 2 Find the maximum *M* and minimum *m* values of the data, and use these values to find *a*, *b*, *c*, and *d*.

[0, 12] scl: 1 by [0, 20] scl: 2

The maximum and minimum hours of daylight are 15.07 and 9.27, respectively. The amplitude *a* is half of the distance between the extrema.

$$a = \frac{1}{2}(M - m) = \frac{1}{2}(15.07 - 9.27)$$
 or 2.9

The vertical shift *d* is the average of the maximum and minimum data values.

 $d = \frac{1}{2}(M + m) = \frac{1}{2}(15.07 + 9.27)$ or 12.17

A sinusoid completes half of a period in the time it takes to go from its maximum to its minimum value. One period is twice this time.

Period = $2(x_{max} - x_{min}) = 2(12 - 6)$ or 12 x_{max} = December 15 or month 12 and x_{min} = June 15 or month 6

Because the period equals $\frac{2\pi}{|b|}$, you can write $|b| = \frac{2\pi}{\text{Period}}$. Therefore, $|b| = \frac{2\pi}{12}$, or $\frac{\pi}{6}$.

The maximum data value occurs when x = 6. Since $y = \cos x$ attains its first maximum when x = 0, we must apply a phase shift of 6 - 0 or 6 units. Use this value to find c.

Phase shift = $-\frac{c}{|b|}$ Phase shift formula $6 = -\frac{c}{\frac{\pi}{6}}$ Phase shift = 6 and $|b| = \frac{\pi}{6}$ $c = -\pi$ Solve for c.

Step 3 Write the function using the values for *a*, *b*, *c*, and *d*. Use $b = \frac{\pi}{6}$.

 $y = 2.9 \cos\left(\frac{\pi}{6}x - \pi\right) + 12.17$ is one model for the hours of daylight

Graph the function and scatter plot in the same viewing window, as in Figure 4.4.1. To find the number of hours of daylight on September 30, evaluate the model for x = 9.5.

$$y = 2.9 \cos\left(\frac{\pi}{6}(9.5) - \pi\right) + 12.17$$
 or about 11.42 hours of daylight

GuidedPractice

METEOROLOGY The average monthly temperatures for Seattle, Washington, are shown.

Month	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sept	Oct	Nov	Dec
Temp. (°)	41	44	47	50	56	61	65	66	61	54	46	42

7A. Write a function that models the monthly temperatures, using x = 1 to represent January.

7B. According to your model, what is Seattle's average monthly temperature in February?

Exercises

 \checkmark

Describe how the graphs of f(x) and g(x) are related. Then find the amplitude of g(x), and sketch two periods of both functions on the same coordinate axes. (Examples 1 and 2)

1. $f(x) = \sin x$	2. $f(x) = \cos x$
$g(x) = \frac{1}{2}\sin x$	$g(x) = -\frac{1}{3}\cos x$
3. $f(x) = \cos x$	4. $f(x) = \sin x$
$g(x) = 6\cos x$	$g(x) = -8\sin x$

Describe how the graphs of f(x) and g(x) are related. Then find the period of g(x), and sketch at least one period of both functions on the same coordinate axes. (Example 3)

5. $f(x) = \sin x$	6. $f(x) = \cos x$
$g(x) = \sin 4x$	$g(x) = \cos 2x$
7. $f(x) = \cos x$	8. $f(x) = \sin x$
$g(x) = \cos\frac{1}{5}x$	$g(x) = \sin\frac{1}{4}x$

9. VOICES The contralto vocal type includes the deepest female singing voice. Some contraltos can sing as low as the E below middle C (E3), which has a frequency of 165 hertz. Write an equation for a sine function that models the initial behavior of the sound wave associated with E3 having an amplitude of 0.15. (Example 4)

Write a sine function that can be used to model the initial behavior of a sound wave with the frequency and amplitude given. (Example 4)

10. $f = 440, a = 0.3$	11. <i>f</i> = 932, <i>a</i> = 0.25
12. $f = 1245, a = 0.12$	13. $f = 623, a = 0.2$

State the amplitude, period, frequency, phase shift, and vertical shift of each function. Then graph two periods of the function. (Examples 5 and 6)

14. $y = 3 \sin\left(x - \frac{\pi}{4}\right)$	15. $y = \cos\left(\frac{x}{3} + \frac{\pi}{2}\right)$
16. $y = 0.25 \cos x + 3$	17. $y = \sin 3x - 2$
18. $y = \cos\left(x - \frac{3\pi}{2}\right) - 1$	19. $y = \sin\left(x + \frac{5\pi}{6}\right) + 4$

20. VACATIONS The average number of reservations *R* that a vacation resort has at the beginning of each month is shown. (Example 7)

Month	R	Month	R
Jan	200	May	121
Feb	173	Jun	175
Mar	113	Jul	198
Apr	87	Aug	168

- **a.** Write an equation of a sinusoidal function that models the average number of reservations using x = 1 to represent January.
- **b.** According to your model, approximately how many reservations can the resort anticipate in November?

21. TIDES The table shown below provides data for the first high and low tides of the day for a certain bay during one day in June. (Example 7)

Tide	Height (ft)	Time
first high tide	12.95	4:25 а.м.
first low tide	2.02	10:55 а.м.

- **a.** Determine the amplitude, period, phase shift, and vertical shift of a sinusoidal function that models the height of the tide. Let *x* represent the number of hours that the high or low tide occurred after midnight.
- **b.** Write a sinusoidal function that models the data.
- **c.** According to your model, what was the height of the tide at 8:45 P.M. that night?
- **22.** METEOROLOGY The average monthly temperatures for Boston, Massachusetts are shown. (Example 7)

Month	Temp. (°F)	Month	Temp. (°F)
Jan	29	Jul	74
Feb	30	Aug	72
Mar	39	Sept	65
Apr	48	Oct	55
May	58	Nov	45
Jun	68	Dec	34

- **a.** Determine the amplitude, period, phase shift, and vertical shift of a sinusoidal function that models the monthly temperatures using x = 1 to represent January.
- **b.** Write an equation of a sinusoidal function that models the monthly temperatures.
- **c.** According to your model, what is Boston's average temperature in August?

GRAPHING CALCULATOR Find the values of *x* in the interval $-\pi < x < \pi$ that make each equation or inequality true. (*Hint:* Use the intersection function.)

23.	$-\sin x = \cos x$	24.	$\sin x - \cos x = 1$
25.	$\sin x + \cos x = 0$	26.	$\cos x \le \sin x$
27.	$\sin x \cos x > 1$	28.	$\sin x \cos x \le 0$

CAROUSELS A wooden horse on a carousel moves up and down as the carousel spins. When the ride ends, the horse usually stops in a vertical position different from where it started. The position *y* of the horse after *t* seconds can be modeled by $y = 1.5 \sin (2t + c)$, where the phase shift *c* must be continuously adjusted to compensate for the different starting positions. If during one ride the horse reached a maximum height after $\frac{7\pi}{12}$ seconds, find the equation that models the horse's position.

30. AMUSEMENT PARKS The position *y* in feet of a passenger cart relative to the center of a Ferris wheel over *t* seconds is shown below.



Side view of Ferris wheel over time interval [0, 5.5]



- **a.** Find the time *t* that it takes for the cart to return to y = 0 during its initial spin.
- **b.** Find the period of the Ferris wheel.
- **c.** Sketch the graph representing the position of the passenger cart over one period.
- **d.** Write a sinusoidal function that models the position of the passenger cart as a function of time *t*.

Write an equation that corresponds to each graph.



Write a sinusoidal function with the given period and amplitude that passes through the given point.

- **35.** period: π ; amplitude: 5; point: $\left(\frac{\pi}{6}, \frac{5}{2}\right)$
- **36.** period: 4π ; amplitude: 2; point: (π , 2)
- **37.** period: $\frac{\pi}{2}$; amplitude: 1.5; point: $\left(\frac{\pi}{2}, \frac{3}{2}\right)$
- **38.** period: 3π ; amplitude: 0.5; point: $\left(\pi, \frac{\sqrt{3}}{4}\right)$

- **39. MULTIPLE REPRESENTATIONS** In this problem, you will investigate the change in the graph of a sinusoidal function of the form $y = \sin x$ or $y = \cos x$ when multiplied by a polynomial function.
 - **a. GRAPHICAL** Use a graphing calculator to sketch the graphs of y = 2x, y = -2x, and $y = 2x \cos x$ on the same coordinate plane, on the interval [-20, 20].
 - **b. VERBAL** Describe the behavior of the graph of $y = 2x \cos x$ in relation to the graphs of y = 2x and y = -2x.
 - **c. GRAPHICAL** Use a graphing calculator to sketch the graphs of $y = x^2$, $y = -x^2$, and $y = x^2 \sin x$ on the same coordinate plane, on the interval [-20, 20].
 - **d. VERBAL** Describe the behavior of the graph of $y = x^2 \sin x$ in relation to the graphs of $y = x^2$ and $y = -x^2$.
 - **e. ANALYTICAL** Make a conjecture as to the behavior of the graph of a sinusoidal function of the form $y = \sin x$ or $y = \cos x$ when multiplied by polynomial function of the form y = f(x).

H.O.T. Problems Use Higher-Order Thinking Skills

40. CHALLENGE Without graphing, find the exact coordinates of the first maximum point to the right of the *y*-axis for $y = 4 \sin\left(\frac{2}{3}x - \frac{\pi}{9}\right)$.

REASONING Determine whether each statement is *true* or *false*. Explain your reasoning.

- **41.** Every sine function of the form $y = a \sin(bx + c) + d \tan(a)$ also be written as a cosine function of the form $y = a \cos(bx + c) + d$.
- **42.** The period of $f(x) = \cos 8x$ is equal to four times the period of $g(x) = \cos 2x$.
- **43 CHALLENGE** How many zeros does $y = \cos 1500x$ have on the interval $0 \le x \le 2\pi$?
- **44. PROOF** Prove the phase shift formula.
- **45.** WRITING IN MATH The Power Tower ride in Sandusky, Ohio, is shown below. Along the side of each tower is a string of lights that send a continuous pulse of light up and down each tower at a constant rate. Explain why the distance *d* of this light from the ground over time *t* cannot be represented by a sinusoidal function.



connectED.mcgraw-hill.com

26

The given point lies on the terminal side of an angle θ in standard position. Find the values of the six trigonometric functions of θ . (Lesson 4-3)

46. (-4, 4) **47.** (8, -2) **48.** (-5, -9) **49.** (4, 5)

52. $-\frac{\pi}{4}$

Write each degree measure in radians as a multiple of π and each radian measure in degrees. (Lesson 4-2)

51. −420°

50. 25°

54. SCIENCE Radiocarbon dating is a method of estimating the age of an organic material by calculating the amount of carbon-14 present in the material. The age of a material can be calculated using $A = t \cdot \frac{\ln R}{-0.693}$, where *A* is the age of the object in years, *t* is the half-life of carbon-14 or 5700 years, and *R* is the ratio of the amount of carbon-14 in the sample to the amount of carbon-14 in living tissue. (Lesson 3-4)

- **a.** A sample of organic material contains 0.000076 gram of carbon-14. A living sample of the same material contains 0.00038 gram. About how old is the sample?
- **b.** A specific sample is at least 20,000 years old. What is the maximum percent of carbon-14 remaining in the sample?

State the number of possible real zeros and turning points of each function. Then determine all of the real zeros by factoring. (Lesson 2-2)

55.	$f(x) = x^3 + 2x^2 - 8x$	56.	$f(x) = x^4 - 10x^2 + 9$
57.	$f(x) = x^5 + 2x^4 - 4x^3 - 8x^2$	58.	$f(x) = x^4 - 1$

Determine whether *f* has an inverse function. If it does, find the inverse function and state any restrictions on its domain. (Lesson 1-7)

59. f(x) = -x - 2 **60.** $f(x) = \frac{1}{x + 4}$ **61.** $f(x) = (x - 3)^2 - 7$ **62.** $f(x) = \frac{1}{(x - 1)^2}$

Skills Review for Standardized Tests 63. SAT/ACT If $x + y = 90^{\circ}$ and x and y are both nonnegative angles, what is equal to $\frac{\cos x}{\sin y}$? **A** 0 D 1.5 $\frac{1}{2}$ B E Cannot be determined from **C** 1 the information given. **64. REVIEW** If $\tan x = \frac{10}{24}$ in the figure below, what are $\sin x$ and $\cos x$? **F** sin $x = \frac{26}{10}$ and cos $x = \frac{24}{26}$ **G** sin $x = \frac{10}{26}$ and cos $x = \frac{24}{26}$ **H** sin $x = \frac{26}{10}$ and cos $x = \frac{26}{24}$ J sin $x = \frac{10}{26}$ and cos $x = \frac{26}{24}$ 266 | Lesson 4-4 | Graphing Sine and Cosine Functions

65. Identify the equation represented by the graph.



sin θ ? F $-\frac{15}{8}$ H $-\frac{15}{17}$

G
$$-\frac{17}{15}$$
 J $-\frac{8}{15}$

53. $\frac{8\pi}{3}$

Graphing Technology Lab Sums and Differences of Sinusoids



Objective

TechnologyTip

make a graph disappear.

Hiding Graphs Scroll to the equals sign and select enter to

 Graph and examine the periods of sums and differences of sinusoids. The graphs of the sums and differences of two sinusoids will often have different periods than the graphs of the original functions.

Activity 1 Sum of Sinusoids



Determine a common interval on which both $f(x) = 2 \sin 3x$ and $g(x) = 4 \cos \frac{x}{2}$ complete a

whole number of cycles. Then graph h(x) = f(x) + g(x), and identify the period of the function.

Step 1 Enter f(x) for Y1 and g(x) for Y2. Then adjust the window until each graph completes one or more whole cycles on the same interval. One interval on which this occurs is $[0, 4\pi]$. On this interval, g(x) completes one whole cycle and f(x) completes six whole cycles.



Step 2 To graph h(x) as Y3, under the VARS menu, select Y-VARS, function, and Y1 to enter Y1. Then press + and select Y-VARS, function, and Y2 to enter Y2.

Step 3 Graph f(x), g(x) and h(x) on the same screen. To make the graph of h(x) stand out, scroll to the left of the equals sign next to Y3, and press **ENTER**. Then graph the functions using the same window as above.



Step 4 By adjusting the x-axis from $[0, 4\pi]$ to $[0, 8\pi]$ to observe the full pattern of h(x), we can see that the period of the sum of the two sinusoids is 4π .



 $[0, 4\pi]$ scl: π by [-6, 6] scl: 1

$\begin{array}{c} y = h(x) \\ y =$

 $[0, 8\pi]$ scl: 2π by [-6, 6] scl: 1

Exercises

Determine a common interval on which both f(x) and g(x) complete a whole number of cycles. Then graph a(x) = f(x) + g(x) and b(x) = f(x) - g(x), and identify the period of the function.

- **1.** $f(x) = 4 \sin 2x$ $g(x) = -2 \cos 3x$ **2.** $f(x) = \sin 8x$ $g(x) = \cos 6x$ **3.** $f(x) = 3 \sin (x - \pi)$ $g(x) = -2 \cos 2x$ **4.** $f(x) = \frac{1}{2} \sin 4x$ $g(x) = 2 \sin \left(x - \frac{\pi}{2}\right)$ **5.** $f(x) = \frac{1}{4} \cos \frac{x}{2}$ $g(x) = -2 \cos \left(x - \frac{\pi}{2}\right)$ **6.** $f(x) = -\frac{1}{2} \sin 2x$ $g(x) = 3 \cos 2x$
- **7.** MAKE A CONJECTURE Explain how you can use the periods of two sinusoids to find the period of the sum or difference of the two sinusoids.

connectED.mcgraw-hill.com 267

Mid-Chapter Quiz

Lessons 4-1 through 4-4

Find the exact values of the six trigonometric functions of θ . (Lesson 4-1)



Find the value of *x*. Round to the nearest tenth if necessary. (Lesson. 4-1)



- **5. SHADOWS** A pine tree casts a shadow that is 7.9 feet long when the Sun is 80° above the horizon. (Lesson 4-1)
 - **a.** Find the height of the tree.
 - **b.** Later that same day, a person 6 feet tall casts a shadow 6.7 feet long. At what angle is the Sun above the horizon?

Find the measure of angle θ . Round to the nearest degree if necessary. (Lesson 4-1)



Identify all angles that are coterminal with the given angle. Then find and draw one positive and one negative angle coterminal with the

9. $\frac{3\pi}{10}$ **10.** -22°

given angle. (Lesson 4-2)

11. MULTIPLE CHOICE Find the approximate area of the shaded region. (Lesson 4-2)



 TRAVEL A car is traveling at a speed of 55 miles per hour on tires that measure 2.6 feet in diameter. Find the approximate angular speed of the tires in radians per minute. (Lesson 4-2)

Sketch each angle. Then find its reference angle. (Lesson 4-3)

13. 175° **14.**
$$\frac{21\pi}{13}$$

Find the exact value of each expression. If undefined, write *undefined.* (Lesson 4-3)

15. cos 315° **16.** sec
$$\frac{3\pi}{2}$$

17.
$$\sin \frac{5\pi}{3}$$
 18. $\tan \frac{5\pi}{6}$

Find the exact values of the five remaining trigonometric functions of θ . (Lesson 4-3)

19.
$$\cos \theta = -\frac{2}{5}$$
, where $\sin \theta < 0$ and $\tan \theta > 0$
20. $\cot \theta = \frac{4}{2}$, where $\cos \theta > 0$ and $\sin \theta > 0$

State the amplitude, period, frequency, phase shift, and vertical shift of each function. Then graph two full periods of the function. (Lesson 4-4)

21.
$$y = -3 \sin\left(x - \frac{3\pi}{2}\right)$$
 22. $y = 5 \cos 2x - 2$

23. MULTIPLE CHOICE Which of the functions has the same graph as $y = 3 \sin (x - \pi)$? (Lesson 4-4)

F
$$y = 3 \sin (x + \pi)$$

G $y = 3 \cos \left(x - \frac{\pi}{2}\right)$
H $y = -3 \sin (x - \pi)$
J $y = -3 \cos \left(x + \frac{\pi}{2}\right)$

- **24. SPRING** The motion of an object attached to a spring oscillating across its original position of rest can be modeled by $x(t) = A \cos \omega t$, where *A* is the initial displacement of the object from its resting position, ω is a constant dependent on the spring and the mass of the object attached to the spring, and *t* is time measured in seconds. (Lesson 4-4)
 - **a.** Draw a graph for the motion of an object attached to a spring and displaced 4 centimeters where $\omega = 3$.
 - **b.** How long will it take for the object to return to its initial position for the first time?
 - **c.** The constant ω is equal to $\sqrt{\frac{k}{m}}$, where *k* is the spring constant, and *m* is the mass of the object. How would increasing the mass of an object affect the period of its oscillations? Explain your reasoning.
- **25. BUOY** The height above sea level in feet of a signal buoy's transmitter is modeled by $h = a \sin bt + \frac{11}{2}$. In rough waters, the height cycles between 1 and 10 feet, with 4 seconds between cycles. Find the

values of a and b.

Graphing Other Trigonometric Functions

Then	Now	: Why?	
 You analyzed graphs of trigonometric functions. (Lesson 4-4) 	 Graph tangent and reciprocal trigonometric functions. Graph damped trigonometric functions. 	• There are two types of radio transmissions known as amplitude modulation (AM) and frequency modulation (FM). When sound is transmitted by an AM radio station, the amplitude of a sinusoidal wave called the <i>carrier wave</i> is changed to produce sound. The transmission of an FM signal results in a change in the frequency of the carrier wave. You will learn more about the graphs of these waves, known as <i>damped waves,</i> in this lesson.	AM signal



BewVocabulary

damped trigonometric function damping factor damped oscillation damped wave damped harmonic motion

Tangent and Reciprocal Functions In Lesson 4-4, you graphed the sine and cosine 1 functions on the coordinate plane. You can use the same techniques to graph the tangent function and the reciprocal trigonometric functions-cotangent, secant, and cosecant.

Since $\tan x = \frac{\sin x}{\cos x}$, the tangent function is undefined when $\cos x = 0$. Therefore, the tangent function has a *vertical asymptote* whenever $\cos x = 0$. Similarly, the tangent and sine functions each have zeros at integer multiples of π because tan x = 0 when sin x = 0.



The properties of the tangent function are summarized below.



StudyTip

Amplitude The term *amplitude* does not apply to the tangent or cotangent functions because the heights of these functions are infinite. The general form of the tangent function, which is similar to that of the sinusoidal functions, is $y = a \tan (bx + c) + d$, where *a* produces a vertical stretch or compression, *b* affects the period, *c* produces a phase shift, *d* produces a vertical shift and neither *a* or *b* are 0.



Two consecutive vertical asymptotes for $y = \tan x$ are $x = -\frac{\pi}{2}$ and $x = \frac{\pi}{2}$. You can find two consecutive vertical asymptotes for any tangent function of the form $y = a \tan(bx + c) + d$ by solving the equations $bx + c = -\frac{\pi}{2}$ and $bx + c = \frac{\pi}{2}$.

You can sketch the graph of a tangent function by plotting the vertical asymptotes, *x*-intercepts, and points between the asymptotes and *x*-intercepts.

Example 1 Graph Horizontal Dilations of the Tangent Function

Locate the vertical asymptotes, and sketch the graph of $y = \tan 2x$.

The graph of $y = \tan 2x$ is the graph of $y = \tan x$ compressed horizontally. The period is $\frac{\pi}{|2|}$ or $\frac{\pi}{2}$. Find two consecutive vertical asymptotes.

$bx + c = -\frac{\pi}{2}$	Tangent asymptote equations	$bx + c = \frac{\pi}{2}$
$2x + 0 = -\frac{\pi}{2}$	b = 2, c = 0	$2x + 0 = \frac{\pi}{2}$
$x = -\frac{\pi}{4}$	Simplify.	$x = \frac{\pi}{4}$

Create a table listing key points, including the *x*-intercept, that are located between the two vertical asymptotes at $x = -\frac{\pi}{4}$ and $x = \frac{\pi}{4}$.

Function	Vertical Asymptote	Intermediate Point	<i>x</i> -intercept	Intermediate Point	Vertical Asymptote
$y = \tan x$	$x = -\frac{\pi}{2}$	$\left(-\frac{\pi}{4},-1\right)$	(0, 0)	$\left(\frac{\pi}{4},1\right)$	$x = \frac{\pi}{2}$
$y = \tan 2x$	$x = -\frac{\pi}{4}$	$\left(-\frac{\pi}{8},-1\right)$	(0, 0)	$\left(\frac{\pi}{8}, 1\right)$	$x = \frac{\pi}{4}$

Sketch the curve through the indicated key points for the function. Then sketch one cycle to the left on the interval $\left(-\frac{3\pi}{4}, -\frac{\pi}{4}\right)$ and one cycle to the right on the interval $\left(\frac{\pi}{4}, \frac{3\pi}{4}\right)$.



GuidedPractice

Locate the vertical asymptotes, and sketch the graph of each function.

1A. $y = \tan 4x$

Example 2 Graph Reflections and Translations of the Tangent Function

Locate the vertical asymptotes, and sketch the graph of each function.

a.
$$y = -\tan \frac{x}{2}$$

The graph of $y = -\tan \frac{x}{2}$ is the graph of $y = \tan x$ expanded horizontally and then reflected in the *x*-axis. The period is $\frac{\pi}{\left|\frac{1}{2}\right|}$ or 2π . Find two consecutive vertical asymptotes.

$$\frac{x}{2} + 0 = -\frac{\pi}{2} \qquad b = \frac{1}{2}, c = 0 \qquad \frac{x}{2} + 0 = \frac{\pi}{2}$$
$$x = 2\left(-\frac{\pi}{2}\right) \text{ or } -\pi \qquad \text{Simplify.} \qquad x = 2\left(\frac{\pi}{2}\right) \text{ or } \pi$$

Create a table listing key points, including the x-intercept, that are located between the two vertical asymptotes at $x = -\pi$ and $x = \pi$.

Function	Vertical Asymptote	Intermediate Point	<i>x</i> -intercept	Intermediate Point	Vertical Asymptote
$y = \tan x$	$x = -\frac{\pi}{2}$	$\left(-\frac{\pi}{4},-1\right)$	(0, 0)	$\left(\frac{\pi}{4}, 1\right)$	$x = \frac{\pi}{2}$
$y = -\tan \frac{x}{2}$	$x = -\pi$	$\left(-\frac{\pi}{2},1\right)$	(0, 0)	$\left(\frac{\pi}{2},-1\right)$	$x = \pi$

Sketch the curve through the indicated key points for the function. Then repeat the pattern for one cycle to the left and right of the first curve.



b.
$$y = \tan\left(x - \frac{3\pi}{2}\right)$$

GuidedPractice

2A. $y = \tan\left(2x + \frac{\pi}{2}\right)$

The graph of $y = \tan\left(x - \frac{3\pi}{2}\right)$ is the graph of $y = \tan x$ shifted $\frac{3\pi}{2}$ units to the right. The period is $\frac{\pi}{|1|}$ or π . Find two consecutive vertical asymptotes.

$$x - \frac{3\pi}{2} = -\frac{\pi}{2} \qquad b = 1, c = -\frac{3\pi}{2} \qquad x - \frac{3\pi}{2} = \frac{\pi}{2}$$
$$x = -\frac{\pi}{2} + \frac{3\pi}{2} \text{ or } \pi \qquad \text{Simplify.} \qquad x = \frac{\pi}{2} + \frac{3\pi}{2} \text{ or } 2\pi$$

Function	Vertical Asymptote	Intermediate Point	<i>x</i> -intercept	Intermediate Point	Vertical Asymptote
$y = \tan x$	$x = -\frac{\pi}{2}$	$\left(-\frac{\pi}{4},-1\right)$	(0, 0)	$\left(\frac{\pi}{4}, 1\right)$	$x = \frac{\pi}{2}$
$y = \tan\left(x - \frac{3\pi}{2}\right)$	$x = \pi$	$\left(\frac{5\pi}{4}, -1\right)$	$\left(\frac{3\pi}{2},0\right)$	$\left(\frac{7\pi}{4},1\right)$	$x = 2\pi$

Sketch the curve through the indicated key points for the function. Then sketch one cycle to the left and right of the first curve.



2B.
$$y = -\tan\left(x - \frac{\pi}{6}\right)$$

StudyTip

Alternate Method When graphing a function with only a horizontal translation *c*, you can find the key points by adding *c* to each of the *x*-coordinates of the key points of the parent function. The cotangent function is the reciprocal of the tangent function, and is defined as $\cot x = \frac{\cos x}{\sin x}$. Like the tangent function, the period of a cotangent function of the form $y = a \cot (bx + c) + d \tan be$ found by calculating $\frac{\pi}{|b|}$. Two consecutive vertical asymptotes can be found by solving the equations bx + c = 0 and $bx + c = \pi$. The properties of the cotangent function are summarized below.



You can sketch the graph of a cotangent function using the same techniques that you used to sketch the graph of a tangent function.

TechnologyTip

Graphing a Cotangent Function When using a calculator to graph a cotangent function, enter the reciprocal of tangent, $y = \frac{1}{\tan x}$. Graphing calculators may produce solid lines where the asymptotes occur. Setting the mode to DOT will eliminate the line.

Example 3 Sketch the Graph of a Cotangent Function

Locate the vertical asymptotes, and sketch the graph of $y = \cot \frac{x}{3}$.

The graph of $y = \cot \frac{x}{3}$ is the graph of $y = \cot x$ expanded horizontally. The period is $\frac{\pi}{\left|\frac{1}{3}\right|}$ or 3π . Find two consecutive vertical asymptotes by solving bx + c = 0 and $bx + c = \pi$.

$$\frac{x}{3} + 0 = 0 \qquad b = \frac{1}{3}, c = 0 \qquad \frac{x}{3} + 0 = \pi$$

x = 3(0) or 0 Simplify. x = 3(\pi) or 3\pi

Create a table listing key points, including the *x*-intercept, that are located between the two vertical asymptotes at x = 0 and $x = 3\pi$.

Function	Vertical Asymptote	Intermediate Point	<i>x</i> -intercept	Intermediate Point	Vertical Asymptote
$y = \cot x$	x = 0	$\left(\frac{\pi}{4},1\right)$	$\left(\frac{\pi}{2}, 0\right)$	$\left(\frac{3\pi}{4},-1\right)$	$x = \pi$
$y = \cot \frac{x}{3}$	x = 0	$\left(\frac{3\pi}{4},1\right)$	$\left(\frac{3\pi}{2},0\right)$	$\left(\frac{9\pi}{4},-1\right)$	$x = 3\pi$

Following the same guidelines that you used for the tangent function, sketch the curve through the indicated key points that you found. Then sketch one cycle to the left and right of the first curve.



GuidedPractice

Locate the vertical asymptotes, and sketch the graph of each function.

3A. $y = -\cot 3x$

The reciprocals of the sine and cosine functions are defined as $\csc x = \frac{1}{\sin x}$ and $\sec x = \frac{1}{\cos x'}$ as shown below.



The cosecant function has asymptotes when sin x = 0, which occurs at integer multiples of π . Likewise, the secant function has asymptotes when $\cos x = 0$, located at odd multiples of $\frac{\pi}{2}$. Notice also that the graph of $y = \csc x$ has a relative minimum at each maximum point on the sine curve, and a relative maximum at each minimum point on the sine curve. The same is true for the graphs of $y = \sec x$ and $y = \cos x$.

The properties of the cosecant and secant functions are summarized below.



Like the sinusoidal functions, the period of a secant function of the form $y = a \sec (bx + c) + d$ or cosecant function of the form $y = a \csc (bx + c) + d$ can be found by calculating $\frac{2\pi}{|b|}$. Two vertical asymptotes for the secant function can be found by solving the equations $bx + c = -\frac{\pi}{2}$ and $bx + c = \frac{3\pi}{2}$ and two vertical asymptotes for the cosecant function can be found by solving $bx + c = -\pi$ and $bx + c = \pi$.

TechnologyTip

Graphing Graphing the cosecant and secant functions on a calculator is similar to graphing the cotangent function. Enter the reciprocals of the sine and cosine functions. To sketch the graph of a cosecant or secant function, locate the asymptotes of the function and find the corresponding relative maximum and minimums points.

Example 4 Sketch Graphs of Cosecant and Secant Functions

Locate the vertical asymptotes, and sketch the graph of each function.

a. $y = \csc\left(x + \frac{\pi}{2}\right)$

The graph of $y = \csc\left(x + \frac{\pi}{2}\right)$ is the graph of $y = \csc x$ shifted $\frac{\pi}{2}$ units to the left. The period is $\frac{2\pi}{|1|}$ or 2π . Two vertical asymptotes occur when $bx + c = -\pi$ and $bx + c = \pi$. Therefore, two asymptotes are $x + \frac{\pi}{2} = -\pi$ or $x = -\frac{3\pi}{2}$ and $x + \frac{\pi}{2} = \pi$ or $x = \frac{\pi}{2}$.

Create a table listing key points, including the relative maximum and minimum, that are located between the two vertical asymptotes at $x = -\frac{3\pi}{2}$ and $x = \frac{\pi}{2}$.

Function	Vertical Asymptote	Relative Maximum	Vertical Asymptote	Relative Minimum	Vertical Asymptote
$y = \csc x$	$x = -\pi$	$\left(-\frac{\pi}{2},-1\right)$	x = 0	$\left(\frac{\pi}{2}, 1\right)$	$x = \pi$
$y = \csc\left(x + \frac{\pi}{2}\right)$	$x = -\frac{3\pi}{2}$	$(-\pi, -1)$	$x = -\frac{\pi}{2}$	(<mark>0</mark> , 1)	$x = \frac{\pi}{2}$

Sketch the curve through the indicated key points for the function. Then sketch one cycle to the left and right. The graph is shown in Figure 4.5.1 below.

b. $y = \sec \frac{x}{4}$

The graph of $y = \sec \frac{x}{4}$ is the graph of $y = \sec x$ expanded horizontally. The period is $\frac{2\pi}{\left|\frac{1}{4}\right|}$ or 8π . Two vertical asymptotes occur when $bx + c = -\frac{\pi}{2}$ and $bx + c = \frac{3\pi}{2}$. Therefore, two asymptotes are $\frac{x}{4} + 0 = -\frac{\pi}{2}$ or $x = -2\pi$ and $\frac{x}{4} + 0 = \frac{3\pi}{2}$ or $x = 6\pi$.

Create a table listing key points that are located between the asymptotes at $x = -2\pi$ and $x = 6\pi$.

Function	Vertical Asymptote	Relative Minimum	Vertical Asymptote	Relative Maximum	Vertical Asymptote
$y = \sec x$	$x = -\frac{\pi}{2}$	(<mark>0</mark> , 1)	$x = \frac{\pi}{2}$	(π , −1)	$x = \frac{3\pi}{2}$
$y = \sec \frac{x}{4}$	$x = -2\pi$	(<mark>0</mark> , 1)	$x = 2\pi$	(4 π, −1)	$x = 6\pi$

Sketch the curve through the indicated key points for the function. Then sketch one cycle to the left and right. The graph is shown in Figure 4.5.2 below.



StudyTip

Finding Asymptotes and Key Points You can use the periodic nature of trigonometric graphs to help find asymptotes and key points. In Example 4a, notice that the vertical asymptote $x = -\frac{\pi}{2}$ is equidistant from the calculated asymptotes, $x = -\frac{3\pi}{2}$ and $x = \frac{\pi}{2}$. **2 Damped Trigonometric Functions** When a sinusoidal function is multiplied by another function f(x), the graph of their product oscillates between the graphs of y = f(x) and y = -f(x). When this product reduces the amplitude of the wave of the original sinusoid, it is called **damped oscillation**, and the product of the two functions is known as a **damped trigonometric function**. This change in oscillation can be seen in Figures 4.5.3 and 4.5.4 for the graphs of $y = \sin x$ and $y = 2x \sin x$.

StudyTip

Damped Functions Trigonometric functions that are multiplied by constants do not experience damping. The constant affects the amplitude of the function.

Math HistoryLink

A Canadian, Morawetz studied the scattering of sound and magnetic

waves and later proved results

relating to the nonlinear wave

Cathleen Synge Morawetz

(1923 -)

equation.



A damped trigonometric function is of the form $y = f(x) \sin bx$ or $y = f(x) \cos bx$, where f(x) is the **damping factor**.

Damped oscillation occurs as *x* approaches $\pm \infty$ or as *x* approaches 0 from both directions.

Example 5 Sketch Damped Trigonometric Functions

Identify the damping factor f(x) of each function. Then use a graphing calculator to sketch the graphs of f(x), -f(x), and the given function in the same viewing window. Describe the behavior of the graph.

a. $y = -3x \cos x$

The function $y = -3x \cos x$ is the product of the functions y = -3x and $y = \cos x$, so f(x) = -3x.

The amplitude of the function is decreasing as *x* approaches 0 from both directions.

b. $y = x^2 \sin x$

The function $y = x^2 \sin x$ is the product of the functions $y = x^2$ and $y = \sin x$. Therefore, the damping factor is $f(x) = x^2$.

The amplitude of the function is decreasing as *x* approaches 0 from both directions.

c. $y = 2^x \cos 3x$

The function $y = 2^x \cos 3x$ is the product of the functions $y = 2^x$ and $y = \cos 3x$, so $f(x) = 2^x$.

The amplitude of the function is decreasing as *x* approaches $-\infty$.

GuidedPractice

5A. $y = 5x \sin x$

5B. $y = \frac{1}{x} \cos x$



 $[-4\pi, 4\pi]$ scl: π by [-40, 40] scl: 5



 $[-4\pi, 4\pi]$ scl: π by [-100, 100] scl: 10



```
5C. y = 3^x \sin x
```

When the amplitude of the motion of an object decreases with time due to friction, the motion is called *damped harmonic motion*.



The greater the damping constant *c*, the faster the amplitude approaches 0. The magnitude of *c* depends on the size of the object and the material of which it is composed.

Real-World Example 6 Damped Harmonic Motion

MUSIC A guitar string is plucked at a distance of 0.8 centimeter above its rest position and then released, causing a vibration. The damping constant for the string is 2.1, and the note produced has a frequency of 175 cycles per second.

a. Write a trigonometric function that models the motion of the string.

The maximum displacement of the string occurs when t = 0, so $y = ke^{-ct}\cos \omega t$ can be used to model the motion of the string because the graph of $y = \cos t$ has a *y*-intercept other than 0.

The maximum displacement occurs when the string is plucked 0.8 centimeter. The total displacement is the maximum displacement M minus the minimum displacement m, so

$$k = M - m = 0.8 - 0$$
 or 0.8 cm.

You can use the value of the frequency to find ω .

$$\frac{|\omega|}{2\pi} = 175 \qquad \frac{|\omega|}{2\pi} = \text{frequency}$$

 $|\omega| = 350\pi$ Multiply each side by 2π .

Write a function using the values of k, ω , and c.

 $y = 0.8e^{-2.1t}\cos 350\pi t$ is one model that describes the motion of the string.

b. Determine the amount of time *t* that it takes the string to be damped so that $-0.28 \le y \le 0.28$.

Use a graphing calculator to determine the value of *t* when the graph of $y = 0.8e^{-2.1t}\cos 350\pi t$ is oscillating between y = -0.28 and y = 0.28.

From the graph, you can see that it takes approximately 0.5 second for the graph of $y = 0.8e^{-2.1t}\cos 350\pi t$ to oscillate within the interval $-0.28 \le y \le 0.28$.



[0, 1] scl: 0.5 by [-0.75, 0.75] scl: 0.25

GuidedPractice

- **6. MUSIC** Suppose another string on the guitar was plucked 0.5 centimeter above its rest position with a frequency of 98 cycles per second and a damping constant of 1.7.
 - **A.** Write a trigonometric function that models the motion of the string *y* as a function of time *t*.
 - **B.** Determine the time *t* that it takes the string to be damped so that $-0.15 \le y \le 0.15$.



Real-WorldLink

Each string on a guitar is stretched to a particular length and tautness. These aspects, along with the weight and type of string, cause it to vibrate with a characteristic frequency or pitch called its fundamental frequency, producing the note we hear.

Source: How Stuff Works

 \checkmark

Locate the vertical asymptotes, and sketch the graph of each function. (Examples 1–4)

1. $y = 2 \tan x$ 3. $y = \cot \left(x - \frac{\pi}{6}\right)$ 5. $y = -\frac{1}{4} \cot x$ 7. $y = -2 \tan (6x - \pi)$ 9. $y = \frac{1}{5} \csc 2x$ 10. $y = \csc \left(\frac{4x + \frac{7\pi}{6}}{6}\right)$ 11. $y = \sec (x + \pi)$ 12. $y = -2 \csc 3x$ 13. $y = 4 \sec \left(x - \frac{3\pi}{4}\right)$ 14. $y = \sec \left(\frac{x}{5} + \frac{\pi}{5}\right)$ 15. $y = \frac{3}{2} \csc \left(x - \frac{2\pi}{3}\right)$ 16. $y = -\sec \frac{x}{8}$

Identify the damping factor f(x) of each function. Then use a graphing calculator to sketch the graphs of f(x), -f(x), and the given function in the same viewing window. Describe the behavior of the graph. (Example 5)

17. $y = \frac{3}{5}x \sin x$	18. $y = 4x \cos x$
19. $y = 2x^2 \cos x$	20. $y = \frac{x^3}{2} \sin x$
21. $y = \frac{1}{3}x \sin 2x$	22. $y = (x - 2)^2 \sin x$
23. $y = e^{0.5x} \cos x$	24. $y = 3^x \sin x$
25. $y = x \cos 3x$	26. $y = \ln x \cos x$

27. MECHANICS When the car shown below hit a bump in the road, the shock absorber was compressed 8 inches, released, and then began to vibrate in damped harmonic motion with a frequency of 2.5 cycles per second. The damping constant for the shock absorber is 3. (Example 6)



- **a.** Write a trigonometric function that models the displacement of the shock absorber *y* as a function of time *t*. Let *t* = 0 be the instant the shock absorber is released.
- **b.** Determine the amount of time *t* that it takes for the amplitude of the vibration to decrease to 4 inches.

28. DIVING The end of a diving board is 20.3 centimeters above its resting position at the moment a diver leaves the board. Two seconds later, the board has moved down and up 12 times. The damping constant for the board is 0.901. (Example 6)



- **a.** Write a trigonometric function that models the motion of the diving board *y* as a function of time *t*.
- **b.** Determine the amount of time *t* that it takes the diving board to be damped so that $-0.5 \le y \le 0.5$.

Locate the vertical asymptotes, and sketch the graph of each function.

29. $y = \sec x + 3$	30. $y = \sec\left(x - \frac{\pi}{2}\right) + 4$
31. $y = \csc \frac{x}{3} - 2$	32. $y = \csc\left(3x + \frac{\pi}{6}\right) + 3$
33. $y = \cot(2x + \pi) - 3$	34. $y = \cot\left(\frac{x}{2} + \frac{\pi}{2}\right) - 1$

35 PHOTOGRAPHY Jeff is taking pictures of a hawk that is flying 150 feet above him. The hawk will eventually fly directly over Jeff. Let *d* be the distance Jeff is from the hawk and θ be the angle of elevation to the hawk from Jeff's camera.



- **a.** Write *d* as a function of θ .
- **b.** Graph the function on the interval $0 < \theta < \pi$.
- **c.** Approximately how far away is the hawk from Jeff when the angle of elevation is 45°?
- **36. DISTANCE** A spider is slowly climbing up a wall. Brianna is standing 6 feet away from the wall watching the spider. Let *d* be the distance Brianna is from the spider and *θ* be the angle of elevation to the spider from Brianna.
 - **a.** Write *d* as a function of θ .
 - **b.** Graph the function on the interval $0 < \theta < \frac{\pi}{2}$.
 - **c.** Approximately how far away is the spider from Brianna when the angle of elevation is 32°?



GRAPHING CALCULATOR Find the values of θ on the interval $-\pi < \theta < \pi$ that make each equation true.

- **37.** $\cot \theta = 2 \sec \theta$ **38.** $\sin \theta = \cot \theta$ **39.** $4 \cos \theta = \csc \theta$ **40.** $\tan \frac{\theta}{2} = \sin \theta$ **41.** $\csc \theta = \sec \theta$ **42.** $\tan \theta = \sec \frac{\theta}{2}$
- **43. TENSION** A helicopter is delivering a large mural that is to be displayed in the center of town. Two ropes are used to attach the mural to the helicopter, as shown. The tension *T* on each rope is equal to half the downward force times $\sec \frac{\theta}{2}$.



- **a.** The downward force in newtons equals the mass of the mural times gravity, which is 9.8 newtons per kilogram. If the mass of the mural is 544 kilograms, find the downward force.
- **b.** Write an equation that represents the tension *T* on each rope.
- **c.** Graph the equation from part **b** on the interval [0, 180°].
- d. Suppose the mural is 9.14 meters long and the ideal angle *θ* for tension purposes is a right angle. Determine how much rope is needed to transport the mural and the tension that is being applied to each rope.
- **e.** Suppose you have 12.2 meters of rope to use to transport the mural. Find *θ* and the tension that is being applied to each rope.

Match each function with its graph.



GRAPHING CALCULATOR Graph each pair of functions on the same screen and make a conjecture as to whether they are equivalent for all real numbers. Then use the properties of the functions to verify each conjecture.

48.
$$f(x) = \sec x \cos x; g(x) = 1$$

49. $f(x) = \sec^2 x; g(x) = \tan^2 x + 1$
50. $f(x) = \cos x \csc x; g(x) = \cot x$
51. $f(x) = \frac{1}{\sec \left(x - \frac{\pi}{2}\right)}; g(x) = \sin x$

Write an equation for the given function given the period, phase shift (ps), and vertical shift (vs).

- **52.** function: sec; period: 3π; ps: 0; vs: 2
- **53** function: tan; period: $\frac{\pi}{2}$; ps: $\frac{\pi}{4}$; vs: -1
- **54.** function: csc; period: $\frac{\pi}{4}$; ps: $-\pi$; vs: 0
- **55.** function: cot; period: 3π ; ps: $\frac{\pi}{2}$; vs: 4
- **56.** function: csc; period: $\frac{\pi}{3}$; ps: $-\frac{\pi}{2}$; vs: -3

H.O.T. Problems Use Higher-Order Thinking Skills

57. PROOF Verify that the *y*-intercept for the graph of any function of the form $y = ke^{-ct} \cos \omega t$ is *k*.

REASONING Determine whether each statement is *true* or *false*. Explain your reasoning.

- **58.** If $b \neq 0$, then $y = a + b \sec x$ has extrema of $\pm (a + b)$.
- **59.** If $x = \theta$ is an asymptote of $y = \csc x$, then $x = \theta$ is also an asymptote of $y = \cot x$.
- **60. ERROR ANALYSIS** Mira and Arturo are studying the graph shown. Mira thinks that it is the graph of $y = -\frac{1}{3} \tan 2x$, and Arturo thinks that it is the graph of $y = \frac{1}{3} \cot 2x$. Is either of them correct? Explain your reasoning.



- **61. CHALLENGE** Write a cosecant function and a cotangent function that have the same graphs as *y* = sec *x* and *y* = tan *x* respectively. Check your answers by graphing.
- **62.** WRITING IN MATH A damped trigonometric function oscillates between the positive and negative graphs of the damping factor. Explain why a damped trigonometric function oscillates between the positive and negative graphs of the damping factor and why the amplitude of the function depends on the damping factor.

278 | Lesson 4-5 | Graphing Other Trigonometric Functions

Spiral Review

State the amplitude, period, frequency, phase shift, and vertical shift of each function. Then graph two periods of the function. (Lesson 4-4)

63. $y = 3\sin\left(2x - \frac{\pi}{3}\right) + 10$ **64.** $y = 2\cos\left(3x + \frac{3\pi}{4}\right) - 6$ **65.** $y = \frac{1}{2}\cos\left(4x - \pi\right) + 1$

Find the exact values of the five remaining trigonometric functions of θ . (Lesson 4-3)

66. $\sin \theta = \frac{4}{5}, \cos \theta > 0$ **67.** $\cos \theta = \frac{6\sqrt{37}}{37}, \sin \theta > 0$ **68.** $\tan \theta = \frac{24}{7}, \sin \theta > 0$

69. POPULATION The population of a city 10 years ago was 45,600. Since then, the population has increased at a steady rate each year. If the population is currently 64,800, find the annual rate of growth for this city. (Lesson 3-5)

70. MEDICINE The half-life of a radioactive substance is the amount of time it takes for half of the atoms of the substance to disintegrate. Nuclear medicine technologists use the iodine isotope I-131, with a half-life of 8 days, to check a patient's thyroid function. After ingesting a tablet containing the iodine, the isotopes collect in the patient's thyroid, and a special camera is used to view its function. Suppose a patient ingests a tablet containing 9 microcuries of I-131. To the nearest hour, how long will it be until there are only 2.8 microcuries in the patient's thyroid? (Lesson 3-2)

Factor each polynomial completely using the given factor and long division. (Lesson 2-3)

71. $x^3 + 2x^2 - x - 2; x - 1$	72. $x^3 + x^2 - 16x - 16; x + 4$	73. $x^3 - x^2 - 10x - 8; x + 1$
--	--	---

74. EXERCISE The American College of Sports Medicine recommends that healthy adults exercise at a target level of 60% to 90% of their maximum heart rates. You can estimate your maximum heart rate by subtracting your age from 220. Write a compound inequality that models age *a* and target heart rate *r*. (Lesson 0-5)

Skills Review for Standardized Tests

75. SAT/ACT In the figure, *A* and *D* are the centers of the two circles, which intersect at points *C* and *E*. \overline{CE} is a diameter of circle *D*. If AB = CE = 10, what is *AD*?





 $\mathbf{F} = \frac{\sqrt{3}}{2}$

G $14\sqrt{3}$

77. Which equation is represented by the graph?

A
$$y = \cot \left(\theta + \frac{\pi}{4}\right)$$

B $y = \cot \left(\theta - \frac{\pi}{4}\right)$
C $y = \tan \left(\theta + \frac{\pi}{4}\right)$
D $y = \tan \left(\theta - \frac{\pi}{4}\right)$
78. REVIEW If $\sin \theta = -\frac{1}{2}$ and $\pi < \theta < \frac{3\pi}{2}$, then $\theta = ?$
F $\frac{13\pi}{12}$
H $\frac{5\pi}{4}$
G $\frac{7\pi}{6}$
J $\frac{4\pi}{3}$

Inverse Trigonometric Functions

Then

Now

Why?

- You found and graphed the inverses of relations and functions.
 You found and inverses functions
 - (Lesson 1-7)

NewVocabulary

arcsine function

arccosine function

arctangent function

Evaluate and graph inverse trigonometric functions.
Find compositions of trigonometric

is not one-to-one.

functions.

Inverse trigonometric functions can be used to model the changing horizontal angle of rotation needed for a television camera to follow the motion of a drag-racing vehicle.

Inverse Trigonometric Functions In Lesson 1-7, you

learned that a function has an inverse function if and

only if it is one-to-one, meaning that each y-value of the

function can be matched with no more than one *x*-value. Because the sine function fails the horizontal line test, it



If, however, we restrict the domain of the sine function to the interval $\left[-\frac{\pi}{2}, \frac{\pi}{2}\right]$, the restricted

function *is* one-to-one and takes on all possible range values [-1, 1] of the unrestricted function. It is on this restricted domain that $y = \sin x$ has an inverse function called the *inverse sine function* $y = \sin^{-1} x$. The graph of $y = \sin^{-1} x$ is found by reflecting the graph of the restricted sine function in the line y = x.



Notice that the domain of $y = \sin^{-1} x$ is [-1, 1], and its range is $\left\lfloor -\frac{\pi}{2}, \frac{\pi}{2} \right\rfloor$. Because angles and arcs given on the unit circle have equivalent radian measures, the inverse sine function is sometimes referred to as the arcsine function $y = \arcsin x$.

In Lesson 4-1, you used the inverse relationship between the sine and inverse sine functions to find acute angle measures. From the graphs above, you can see that in general,

$$y = \sin^{-1} x$$
 or $y = \arcsin x$ iff $\sin y = x$, when $-1 \le x \le 1$ and $-\frac{\pi}{2} \le y \le \frac{\pi}{2}$. iff means if and only if.

This means that $\sin^{-1} x$ or arcsin *x* can be interpreted as *the angle (or arc) between* $-\frac{\pi}{2}$ and $\frac{\pi}{2}$ with a sine of *x*. For example, $\sin^{-1} 0.5$ is the angle with a sine of 0.5.

Recall that sin *t* is the *y*-coordinate of the point on the unit circle corresponding to the angle or arc length *t*. Because the range of

the inverse sine function is restricted to $\left[-\frac{\pi}{2}, \frac{\pi}{2}\right]$, the possible angle measures of the inverse sine function are located on the right half of the unit circle, as shown.



You can use the unit circle to find the exact value of some expressions involving $\sin^{-1} x$ or $\arcsin x$.



Notice in Example 1a that while $\sin \frac{5\pi}{6}$ is also $\frac{1}{2}$, $\frac{5\pi}{6}$ is not in the interval $\left[-\frac{\pi}{2}, \frac{\pi}{2}\right]$. Therefore, $\sin^{-1}\frac{1}{2} \neq \frac{5\pi}{6}$.





StudyTip

Principal Values Trigonometric functions with restricted domains are sometimes indicated with capital letters. For example, $y = \sin x$ represents the function $y = \sin x$, where $-\frac{\pi}{2} \le x \le \frac{\pi}{2}$. The values in these restricted domains are often called *principal values*. When restricted to a domain of $[0, \pi]$, the cosine function is one-to-one and takes on all of its possible range values on [-1, 1]. It is on this restricted domain that the cosine function has an inverse function, called the *inverse cosine function* $y = \cos^{-1} x$ or arccosine function $y = \arccos^{-1} x$. The graph of $y = \cos^{-1} x$ is found by reflecting the graph of the restricted cosine function in the line y = x.



Recall that $\cos t$ is the *x*-coordinate of the point on the unit circle corresponding to the angle or arc length *t*. Because the range of $y = \cos^{-1} x$ is restricted to $[0, \pi]$, the values of an inverse cosine function are located on the upper half of the unit circle.



Example 2 Evaluate Inverse Cosine Functions

Find the exact value of each expression, if it exists.

a. $\cos^{-1}\left(-\frac{\sqrt{2}}{2}\right)$

Find a point on the unit circle in the interval $[0, \pi]$ with an *x*-coordinate of $-\frac{\sqrt{2}}{2}$. When $t = \frac{3\pi}{4}$, $\cos t = -\frac{\sqrt{2}}{2}$. Therefore, $\cos^{-1}\left(-\frac{\sqrt{2}}{2}\right) = \frac{3\pi}{4}$.

CHECK If $\cos^{-1}\left(-\frac{\sqrt{2}}{2}\right) = \frac{3\pi}{4}$, then $\cos\frac{3\pi}{4} = -\frac{\sqrt{2}}{2}$.



b. arccos (-2)

Since the domain of the cosine function is [-1, 1] and -2 < -1, there is no angle with a cosine of -2. Therefore, the value of arccos (-2) does not exist.

c. $\cos^{-1} 0$

Find a point on the unit circle in the interval $[0, \pi]$ with an *x*-coordinate of 0. When $t = \frac{\pi}{2}$, cos t = 0.

Therefore, $\cos^{-1} 0 = \frac{\pi}{2}$

CHECK If $\cos^{-1} 0 = \frac{\pi}{2}$, then $\cos \frac{\pi}{2} = 0$.

GuidedPractice

2A. $\cos^{-1}\left(-\frac{\sqrt{3}}{2}\right)$

2B. arccos 2.5



2C. $\cos^{-1}\left(-\frac{1}{2}\right)$

StudyTip

End Behavior of Inverse Tangent Notice that when the graph of the restricted tangent function is reflected in the line y = x, the vertical asymptotes at $x = \pm \frac{\pi}{2}$ become the horizontal asymptotes $y = \pm \frac{\pi}{2}$ of the inverse tangent function. Therefore, $\lim_{x \to -\infty} \tan^{-1} x = -\frac{\pi}{2}$ and $\lim_{x \to \infty} \tan^{-1} x = \frac{\pi}{2}$. When restricted to a domain of $\left(-\frac{\pi}{2}, \frac{\pi}{2}\right)$, the tangent function is one-to-one. It is on this restricted domain that the tangent function has an inverse function called the *inverse tangent function* $y = \tan^{-1} x$ or **arctangent function** $y = \arctan x$. The graph of $y = \tan^{-1} x$ is found by reflecting the graph of the restricted tangent function in the line y = x. Notice that unlike the sine and cosine functions, the domain of the inverse tangent function is $(-\infty, \infty)$.





You can also use the unit circle to find the value of an inverse tangent expression. On the unit circle, $\tan t = \frac{\sin t}{\cos t} \operatorname{or} \frac{y}{x}$. The values of $y = \tan^{-1} x$ will be located on the right half of the unit circle, not including $-\frac{\pi}{2}$ and $\frac{\pi}{2}$, because the tangent function is undefined at those points.



TechnologyTip

Evaluate \tan^{-1} You can also use a graphing calculator to find the angle that has a tangent of $\sqrt{3}$.



Example 3 Evaluate Inverse Tangent Functions

Find the exact value of each expression, if it exists.

a. $\tan^{-1}\sqrt{3}$

Find a point (x, y) on the unit circle in the interval $\left(-\frac{\pi}{2}, \frac{\pi}{2}\right)$ such that $\frac{y}{x} = \sqrt{3}$. When $t = \frac{\pi}{3}$, $\tan t = \frac{\frac{\sqrt{3}}{2}}{\frac{1}{2}}$ or $\sqrt{3}$. Therefore, $\tan^{-1}\sqrt{3} = \frac{\pi}{3}$.

CHECK If $\tan^{-1}\sqrt{3} = \frac{\pi}{3}$, then $\tan\frac{\pi}{3} = \sqrt{3}$.

b. arctan 0

Find a point (*x*, *y*) on the unit circle in the interval $\left(-\frac{\pi}{2}, \frac{\pi}{2}\right)$ such that $\frac{y}{x} = 0$. When t = 0, $\tan t = \frac{0}{1}$ or 0.

Therefore, $\arctan 0 = 0$.

CHECK If $\arctan 0 = 0$, then $\tan 0 = 0$.







While inverse functions for secant, cosecant, and cotangent do exist, these functions are rarely used in computations because the inverse functions for their reciprocals exist. Also, deciding how to restrict the domains of secant, cosecant, and cotangent to obtain arcsecant, arccosecant, and arccotangent is not as apparent. You will explore these functions in Exercise 66.





connectED.mcgraw-hill.com

283



You can sketch the graph of one of the inverse trigonometric functions shown above by rewriting the function in the form $\sin y = x$, $\cos y = x$, or $\tan y = x$, assigning values to y and making a table of values, and then plotting the points and connecting the points with a smooth curve.

Example 4 Sketch Graphs of Inverse Trigonometric Functions

Sketch the graph of $y = \arccos 2x$.

By definition, $y = \arccos 2x$ and $\cos y = 2x$ are equivalent on $0 \le y \le \pi$, so their graphs are the same. Rewrite $\cos y = 2x$ as $x = \frac{1}{2} \cos y$ and assign values to y on the interval $[0, \pi]$ to make a table of values.

y	0	$\frac{\pi}{4}$	$\frac{\pi}{6}$	$\frac{\pi}{2}$	$\frac{5\pi}{6}$	$\frac{3\pi}{4}$	π
$x=\frac{1}{2}\cos y$	$\frac{1}{2}$	$\frac{\sqrt{2}}{4}$	$\frac{\sqrt{3}}{4}$	0	$-\frac{\sqrt{3}}{4}$	$-\frac{\sqrt{2}}{4}$	$-\frac{1}{2}$

Then plot the points (x, y) and connect them with a smooth curve. Notice that this curve has endpoints at $\left(-\frac{1}{2}, \pi\right)$ and $\left(\frac{1}{2}, 0\right)$, indicating that the entire graph of $y = \arccos 2x$ is shown.



GuidedPractice

Sketch the graph of each function.

4A. $y = \arcsin 3x$







Real-WorldLink

In the late 19th century, Thomas Edison began work on a device to record moving images, called the kinetoscope, which would later become the film projector. The earliest copyrighted motion picture is a film of one of Edison's employees sneezing.

Source: The Library of Congress

Real-World Example 5 Use an Inverse Trigonometric Function

MOVIES In a movie theater, a person's viewing angle for watching a movie changes depending on where he or she sits in the theater.

a. Write a function modeling the viewing angle θ for a person in the theater whose eye-level when sitting is 4 feet above ground.

Draw a diagram to find the measure of the viewing angle. Let θ_1 represent the angle formed from eye-level to the bottom of the screen, and let θ_2 represent the angle formed from eye-level to the top of the screen.



So, the viewing angle is $\theta = \theta_2 - \theta_1$. You can use the tangent function to find θ_1 and θ_2 . Because the eye-level of the person when seated is 4 feet above the floor, the distance opposite θ_1 is 8 - 4 feet or 4 feet long.

$$\tan \theta_1 = \frac{4}{d}$$
 opp = 4 and adj = d
 $\theta_1 = \tan^{-1} \frac{4}{d}$ Inverse tangent function

The distance opposite θ_2 is (32 + 8) - 4 feet or 36 feet.

 $\tan \theta_2 = \frac{36}{d} \qquad \text{opp} = 36 \text{ and } \text{adj} = d$ $\theta_2 = \tan^{-1} \frac{36}{d} \qquad \text{Inverse tangent function}$

So, the viewing angle can be modeled by $\theta = \tan^{-1} \frac{36}{d} - \tan^{-1} \frac{4}{d}$.

b. Determine the distance that corresponds to the maximum viewing angle.

The distance at which the maximum viewing angle occurs is the maximum point on the graph. You can use a graphing calculator to find this point.

From the graph, you can see that the maximum viewing angle occurs approximately 12 feet from the screen.



[0, 100] scl: 10 by [0, 60] scl: 5

GuidedPractice

5. TELEVISION Tucas has purchased a new flat-screen television. So that his family will be able to see, he has decided to hang the television on the wall as shown.



- **A.** Write a function modeling the distance *d* of the maximum viewing angle θ for Lucas if his eye level when sitting is 3 feet above ground.
- **B.** Determine the distance that corresponds to the maximum viewing angle.

2 Compositions of Trigonometric Functions In Lesson 1-7, you learned that if *x* is in the domain of f(x) and $f^{-1}(x)$, then

 $f[f^{-1}(x)] = x$ and $f^{-1}[f(x)] = x$.

Because the domains of the trigonometric functions are restricted to obtain the inverse trigonometric functions, the properties do not apply for all values of x.

For example, while sin *x* is defined for all *x*, the domain of sin⁻¹ *x* is [-1, 1]. Therefore, sin (sin⁻¹ *x*) = *x* is only true when $-1 \le x \le 1$. A different restriction applies for the composition sin⁻¹ (sin *x*). Because the domain of sin *x* is restricted to the interval $\left[-\frac{\pi}{2}, \frac{\pi}{2}\right]$, sin⁻¹ (sin *x*) = *x* is only true when $-\frac{\pi}{2} \le x \le \frac{\pi}{2}$.

These domain restrictions are summarized below.

KeyConcept Domain of Compositions of Trigonometric Functions			
$f[f^{-1}(x)] = x$	$f^{-1}[f(x)] = x$		
If $-1 \le x \le 1$, then sin $(\sin^{-1} x) = x$.	If $-\frac{\pi}{2} \le x \le \frac{\pi}{2}$, then $\sin^{-1}(\sin x) = x$.		
If $-1 \le x \le 1$, then $\cos(\cos^{-1} x) = x$.	If $0 \le x \le \pi$, then $\cos^{-1}(\cos x) = x$.		
If $-\infty < x < \infty$, then $\tan(\tan^{-1} x) = x$.	If $-\frac{\pi}{2} < x < \frac{\pi}{2}$, then $\tan^{-1}(\tan x) = x$.		

Example 6 Use Inverse Trigonometric Properties

Find the exact value of each expression, if it exists.

a. $\sin\left[\sin^{-1}\left(-\frac{1}{4}\right)\right]$

The inverse property applies because $-\frac{1}{4}$ lies on the interval [-1, 1].

Therefore, $\sin\left[\sin^{-1}\left(-\frac{1}{4}\right)\right] = -\frac{1}{4}$.

b. arctan $\left(\tan\frac{\pi}{2}\right)$

Because $\tan x$ is not defined when $x = \frac{\pi}{2}$, $\arctan\left(\tan\frac{\pi}{2}\right)$ does not exist.

c. $\arcsin\left(\sin\frac{7\pi}{4}\right)$ Notice that the angle $\frac{7\pi}{4}$ does not lie on the interval $\left[-\frac{\pi}{2}, \frac{\pi}{2}\right]$. However, $\frac{7\pi}{4}$ is coterminal with $\frac{7\pi}{4} - 2\pi$ or $-\frac{\pi}{4}$, which is on the interval $\left[-\frac{\pi}{2}, \frac{\pi}{2}\right]$. $\arcsin\left(\sin\frac{7\pi}{4}\right) = \arcsin\left[\sin\left(-\frac{\pi}{4}\right)\right]$ $\sin\frac{7\pi}{4} = \sin\left(-\frac{\pi}{4}\right)$ $= -\frac{\pi}{4}$ $\operatorname{Since} -\frac{\pi}{2} \le -\frac{\pi}{4} \le \frac{\pi}{2}$, $\arcsin\left(\sin x\right) = x$. Therefore, $\arcsin\left(\sin\frac{7\pi}{4}\right) = -\frac{\pi}{4}$. **GuidedPractice 6A.** $\tan\left(\tan^{-1}\frac{\pi}{3}\right)$ **6B.** $\cos^{-1}\left(\cos\frac{3\pi}{4}\right)$ **6C.** $\arcsin\left(\sin\frac{2\pi}{3}\right)$

WatchOut!

Compositions and Inverses When computing $f^{-1}[f(x)]$ with trigonometric functions, the domain appears to be $(-\infty, \infty)$. However, because the ranges of the inverse functions are restricted, coterminal angles must sometimes be found. You can also evaluate the composition of two different inverse trigonometric functions.



Sometimes the composition of two trigonometric functions reduces to an algebraic expression that does not involve any trigonometric expressions.

StudyTip

Decomposing Algebraic Functions The technique used to convert a trigonometric expression into an algebraic expression can be reversed. Decomposing an algebraic function as the composition of two trigonometric functions is a technique used frequently in calculus.

Example 8 Evaluate Compositions of Trigonometric Functions

Write tan (arcsin *a*) as an algebraic expression of *a* that does not involve trigonometric functions.

Let $u = \arcsin a$, so $\sin u = a$.

Because the domain of the inverse sine function is restricted to Quadrants I and IV, u must lie in Quadrant I or IV. The solution is similar for each quadrant, so we will solve for Quadrant I.

From the Pythagorean Theorem, you can find that the length of the side adjacent to *u* is $\sqrt{1-a^2}$. Now, solve for tan *u*.

$$\tan u = \frac{\text{opp}}{\text{adj}}$$
Tangent function
$$= \frac{a}{\sqrt{1 - a^2}} \text{ or } \frac{a\sqrt{1 - a^2}}{1 - a^2} \quad \text{opp} = a \text{ and adj}$$

$$pp = a$$
 and $adi = \sqrt{1 - a^2}$

So,
$$\tan(\arcsin a) = \frac{a\sqrt{1-a^2}}{1-a^2}$$

GuidedPractice

Write each expression as an algebraic expression of x that does not involve trigonometric functions.

8A. $\sin(\arccos x)$

8B. $\cot [\sin^{-1} x]$



Find the exact value of each expression, if it exists. (Examples 1–3)

1.	$\sin^{-1} 0$	2.	$\arcsin\frac{\sqrt{3}}{2}$
3.	$\arcsin\frac{\sqrt{2}}{2}$	4.	$\sin^{-1}\frac{1}{2}$
5.	$\sin^{-1}\left(-\frac{\sqrt{2}}{2}\right)$	6.	arccos 0
7.	$\cos^{-1}\frac{\sqrt{2}}{2}$	8.	$\arccos(-1)$
9.	$\arccos \frac{\sqrt{3}}{2}$	10.	$\cos^{-1}\frac{1}{2}$
11.	arctan 1	12.	$\arctan(-\sqrt{3})$
13.	$\tan^{-1}\frac{\sqrt{3}}{3}$	14.	$\tan^{-1} 0$

15. ARCHITECTURE The support for a roof is shaped like two right triangles, as shown below. Find *θ*. (Example 3)



16. RESCUE A cruise ship sailed due west 24 miles before turning south. When the cruise ship became disabled and the crew radioed for help, the rescue boat found that the fastest route covered a distance of 48 miles. Find the angle θ at which the rescue boat should travel to aid the cruise ship. (Example 3)



Sketch the graph of each function. (Example 4)

17. $y = \arcsin x$	18. $y = \sin^{-1} 2x$
19. $y = \sin^{-1}(x+3)$	20. $y = \arcsin x - 3$
21. $y = \arccos x$	22. $y = \cos^{-1} 3x$
23. $y = \arctan x$	24. $y = \tan^{-1} 3x$
25. $y = \tan^{-1}(x+1)$	26. $v = \arctan x - 1$

27 DRAG RACE A television camera is filming a drag race. The camera rotates as the vehicles move past it. The camera is 30 meters away from the track. Consider θ and *x* as shown in the figure. (Example 5)



- **a.** Write θ as a function of *x*.
- **b.** Find θ when x = 6 meters and x = 14 meters.
- **28. SPORTS** Steve and Ravi want to project a pro soccer game on the side of their apartment building. They have placed a projector on a table that stands 5 feet above the ground and have hung a 12-foot-tall screen that is 10 feet above the ground. (Example 5)



- **a.** Write a function expressing θ in terms of distance *d*.
- **b.** Use a graphing calculator to determine the distance for the maximum projecting angle.

Find the exact value of each expression, if it exists. (Examples 6 and 7)

29. $\sin\left(\sin^{-1}\frac{3}{4}\right)$ **30.** $\sin^{-1}\left(\sin\frac{\pi}{2}\right)$
31. $\cos\left(\cos^{-1}\frac{2}{9}\right)$ **32.** $\cos^{-1}(\cos\pi)$
33. $\tan\left(\tan^{-1}\frac{\pi}{4}\right)$ **34.** $\tan^{-1}\left(\tan\frac{\pi}{3}\right)$
35. $\cos(\tan^{-1}1)$ **36.** $\sin^{-1}\left(\cos\frac{\pi}{2}\right)$
37. $\sin\left(2\cos^{-1}\frac{\sqrt{2}}{2}\right)$ **38.** $\sin(\tan^{-1}1 - \sin^{-1}1)$
39. $\cos(\tan^{-1}1 - \sin^{-1}1)$ **40.** $\cos\left(\cos^{-1}0 + \sin^{-1}\frac{1}{2}\right)$

Write each trigonometric expression as an algebraic expression of *x*. (Example 8)

(41)	tan (arccos x)	42.	$\csc(\cos^{-1}x)$
43.	$\sin\left(\cos^{-1}x\right)$	44.	$\cos(\arcsin x)$
45.	$\csc(\sin^{-1}x)$	46.	sec (arcsin <i>x</i>)
47.	$\cot(\arccos x)$	48.	$\cot(\arcsin x)$

0

Describe how the graphs of g(x) and f(x) are related.

49. $f(x) = \sin^{-1} x$ and $g(x) = \sin^{-1} (x - 1) - 2$

- **50.** $f(x) = \arctan x$ and $g(x) = \arctan 0.5x 3$
- **51.** $f(x) = \cos^{-1} x$ and $g(x) = 3 (\cos^{-1} x 2)$
- **52.** $f(x) = \arcsin x$ and $g(x) = \frac{1}{2} \arcsin (x + 2)$
- **53.** $f(x) = \arccos x$ and $g(x) = 5 + \arccos 2x$
- **54.** $f(x) = \tan^{-1} x$ and $g(x) = \tan^{-1} 3x 4$
- **55. SAND** When piling sand, the angle formed between the pile and the ground remains fairly consistent and is called the *angle of repose*. Suppose Jade creates a pile of sand at the beach that is 3 feet in diameter and 1.1 feet high.



- **a.** What is the angle of repose?
- **b.** If the angle of repose remains constant, how many feet in diameter would a pile need to be to reach a height of 4 feet?

Give the domain and range of each composite function. Then use your graphing calculator to sketch its graph.

56. $y = \cos(\tan^{-1} x)$	57. $y = \sin(\cos^{-1} x)$
58. <i>y</i> = arctan (sin <i>x</i>)	59. $y = \sin^{-1} (\cos x)$
60. $y = \cos(\arcsin x)$	61. $y = \tan(\arccos x)$

- **62. INVERSES** The arcsecant function is graphed by restricting the domain of the secant function to the intervals $\left[0, \frac{\pi}{2}\right)$ and $\left(\frac{\pi}{2}, \pi\right]$, and the arccosecant function is graphed by restricting the domain of the cosecant function to the intervals $\left[-\frac{\pi}{2}, 0\right)$ and $\left(0, \frac{\pi}{2}\right]$.
 - **a.** State the domain and range of each function.
 - **b.** Sketch the graph of each function.
 - **c.** Explain why a restriction on the domain of the secant and cosecant functions is necessary in order to graph the inverse functions.

Write each algebraic expression as a trigonometric function of an inverse trigonometric function of *x*.

63.
$$\frac{x}{\sqrt{1-x^2}}$$

64.
$$\frac{\sqrt{1-x^2}}{x}$$

- **65. MULTIPLE REPRESENTATIONS** In this problem, you will explore the graphs of compositions of trigonometric functions.
 - **a. ANALYTICAL** Consider $f(x) = \sin x$ and $f^{-1}(x) = \arcsin x$. Describe the domain and range of $f \circ f^{-1}$ and $f^{-1} \circ f$.
 - **b. GRAPHICAL** Create a table of several values for each composite function on the interval [-2, 2]. Then use the table to sketch the graphs of $f \circ f^{-1}$ and $f^{-1} \circ f$. Use a graphing calculator to check your graphs.
 - **c. ANALYTICAL** Consider $g(x) = \cos x$ and $g^{-1}(x) = \arccos x$. Describe the domain and range of $g \circ g^{-1}$ and $g^{-1} \circ g$ and make a conjecture as to what the graphs of $g \circ g^{-1}$ and $g^{-1} \circ g$ will look like. Explain your reasoning.
 - **d. GRAPHICAL** Sketch the graphs of $g \circ g^{-1}$ and $g^{-1} \circ g$. Use a graphing calculator to check your graphs.
 - e. VERBAL Make a conjecture as to what the graphs of the two possible compositions of the tangent and arctangent functions will look like. Explain your reasoning. Then check your conjecture using a graphing calculator.

H.O.T. Problems Use Higher-Order Thinking Skills

- **66. ERROR ANALYSIS** Alisa and Trey are discussing inverse trigonometric functions. Because $\tan x = \frac{\sin x}{\cos x}$, Alisa conjectures that $\tan^{-1} x = \frac{\sin^{-1} x}{\cos^{-1} x}$. Trey disagrees. Is either of them correct? Explain.
- **67.** CHALLENGE Use the graphs of $y = \sin^{-1} x$ and $y = \cos^{-1} x$ to find the value of $\sin^{-1} x + \cos^{-1} x$ on the interval [-1, 1]. Explain your reasoning.
- **68. REASONING** Determine whether the following statement is *true* or *false*: If $\cos \frac{7\pi}{4} = \frac{\sqrt{2}}{2}$, then $\cos^{-1} \frac{\sqrt{2}}{2} = \frac{7\pi}{4}$. Explain your reasoning.

REASONING Determine whether each function is *odd*, *even*, or *neither*. Justify your answer.

69. $y = \sin^{-1} x$

70.
$$y = \cos^{-1} x$$

- **71.** $y = \tan^{-1} x$
- **72.** WRITING IN MATH Explain how the restrictions on the sine, cosine, and tangent functions dictate the domain and range of their inverse functions.


Spiral Review

Locate the vertical asymptotes, and sketch the graph of each function. (Lesson 4-5)

73. $y = 3 \tan \theta$

75.
$$y = 3 \csc \frac{1}{2} \theta$$

76. WAVES A leaf floats on the water bobbing up and down. The distance between its highest and lowest points is 4 centimeters. It moves from its highest point down to its lowest point and back to its highest point every 10 seconds. Write a cosine function that models the movement of the leaf in relationship to the equilibrium point. (Lesson 4-4)

74. $y = \cot 5\theta$

Find the value of *x*. Round to the nearest tenth, if necessary. (Lesson 4-1)





For each pair of functions, find $[f \circ g](x)$, $[g \circ f](x)$, and $[f \circ g](4)$. (Lesson 1-6)

80. $f(x) = x^2 + 3x - 6$	81. $f(x) = 6 - 5x$	82. $f(x) = \sqrt{x+3}$
g(x) = 4x + 1	$g(x) = \frac{1}{x}$	$g(x) = x^2 + 1$

83. EDUCATION Todd has answered 11 of his last 20 daily quiz questions correctly. His baseball coach told him that he must raise his average to at least 70% if he wants to play in the season opener. Todd vows to study diligently and answer all of the daily quiz questions correctly in the future. How many consecutive daily quiz questions must he answer correctly to raise his average to 70%? (Lesson 0-8)

Skills Review for Standardized Tests

84. SAT/ACT To the nearest degree, what is the angle of depression θ between the shallow end and the deep end of the swimming pool?



85. Which of the following represents the exact value of $\sin\left(\tan^{-1}\frac{1}{2}\right)$?

F
$$-\frac{2\sqrt{5}}{5}$$
 H $\frac{\sqrt{5}}{5}$
G $-\frac{\sqrt{5}}{5}$ **J** $\frac{2\sqrt{5}}{5}$

(

86. REVIEW The hypotenuse of a right triangle is 67 inches. If one of the angles has a measure of 47°, what is the length of the shortest leg of the triangle?

Α	45.7 in.	С	62.5 in.
В	49.0 in.	D	71.8 in.

87. REVIEW Two trucks, *A* and *B*, start from the intersection *C* of two straight roads at the same time. Truck *A* is traveling twice as fast as truck *B* and after 4 hours, the two trucks are 350 miles apart. Find the approximate speed of truck *B* in miles per hour.



The Law of Sines and the Law of Cosines

Then	Now	: Why?	
 You solved right triangles using trigonometric functions. (Lesson 4-1) 	 Solve oblique triangles by using the Law of Sines or the Law of Cosines. Find areas of oblique triangles. 	Triangulation is the process of finding the coordinates of a point and the distance to that point by calculating the length of one side of a triangle, given the measurements of the angles and sides of the triangle formed by that point and two other known reference points. Weather spotters can use triangulation to determine the location of a tornado.	

NewVocabulary
 oblique triangles
 Law of Sines

Law of Sines ambiguous case Law of Cosines Heron's Formula **Solve Oblique Triangles** In Lesson 4-1, you used trigonometric functions to solve *right* triangles. In this lesson, you will solve **oblique triangles**—triangles that are not right triangles.

You can apply the **Law of Sines** to solve an oblique triangle if you know the measures of two angles and a nonincluded side (AAS), two angles and the included side (ASA), or two sides and a nonincluded angle (SSA).

KeyConcept Law of Sines

If $\triangle ABC$ has side lengths *a*, *b*, and *c* representing the lengths of the sides opposite the angles with measures

A, B, and C, then
$$\frac{\sin A}{a} = \frac{\sin B}{b} = \frac{\sin C}{c}$$
.



20

You will derive the Law of Sines in Exercise 69.

Example 1 Apply the Law of Sines (AAS)

Solve $\triangle ABC$. Round side lengths to the nearest tenth and angle measures to the nearest degree.

Because two angles are given, $C = 180^{\circ} - (103^{\circ} + 35^{\circ})$ or 42° .

Use the Law of Sines to find *a* and *c*.



Therefore, $a \approx 11.8$, $c \approx 13.7$, and $\angle C = 42^{\circ}$.

GuidedPractice Solve each triangle. Round side lengths to the nearest tenth and angle measures to the nearest degree.



1A.





Real-WorldLink

To gain an improved understanding of the atmosphere, land surface changes, and ecosystem processes, NASA uses a series of satellites as part of its Earth Observing System (EOS) to study the air, land, and water on Earth.

Source: NASA

Real-World Example 2 Apply the Law of Sines (ASA)

SATELLITES An Earth-orbiting satellite is passing between the Oak Ridge Laboratory in Tennessee and the Langley Research Center in Virginia, which are 446 miles apart. If the angles of elevation to the satellite from the Oak Ridge and Langley facilities are 58° and 72°, respectively, how far is the satellite from each station?



Because two angles are given, $C = 180^{\circ} - (58^{\circ} + 72^{\circ})$ or 50°. Use the Law of Sines to find the distance to the satellite from each station.

$\frac{\sin C}{c} = \frac{\sin B}{b}$	Law of Sines	$\frac{\sin C}{c} = \frac{\sin A}{a}$
$\frac{\sin 50^\circ}{446} = \frac{\sin 72^\circ}{b}$	Substitution	$\frac{\sin 50^\circ}{446} = \frac{\sin 58^\circ}{a}$
$b \sin 50^\circ = 446 \sin 72^\circ$	Multiply.	$a\sin 50^\circ = 446\sin 58^\circ$
$b = \frac{446\sin 72^\circ}{\sin 50^\circ}$	Divide.	$a = \frac{446\sin 58^\circ}{\sin 50^\circ}$
$b \approx 553.72$	Use a calculator.	$a \approx 493.74$

So, the satellite is about 554 miles from Oak Ridge and about 494 miles from Langley.

GuidedPractice

2. SHIPPING Two ships are 250 feet apart and traveling to the same port as shown. Find the distance from the port to each ship.



From geometry, you know that the measures of two sides and a nonincluded angle (SSA) do not necessarily define a unique triangle. Consider the angle and side measures given in the figures below.



In general, given the measures of two sides and a nonincluded angle, one of the following will be true: (1) no triangle exists, (2) exactly one triangle exists, or (3) two triangles exist. In other words, when solving an oblique triangle for this **ambiguous case**, there may be no solution, one solution, or two solutions.

StudyTip

Alternative Representations The Law of Sines can also be written in reciprocal form as





To solve an ambiguous case oblique triangle, first determine the number of possible solutions. If the triangle has one or two solutions, use the Law of Sines to find them.

Example 3 The Ambiguous Case—One or No Solution

Find all solutions for the given triangle, if possible. If no solution exists, write no solution. Round side lengths to the nearest tenth and angle measures to the nearest degree.

a. $a = 15, c = 12, A = 94^{\circ}$

Notice that *A* is obtuse and a > c because 15 > 12. Therefore, one solution exists. Apply the Law of Sines to find C.

$$\frac{\sin C}{12} = \frac{\sin 94^\circ}{15}$$
$$\sin C = \frac{12\sin 94^\circ}{15}$$

Multiply each side by 12. Definition of sin⁻¹

Law of Sines



Because two angles are now known, $B \approx 180^{\circ} - (94^{\circ} + 53^{\circ})$ or about 33°. Apply the Law of Sines to find *b*. Choose the ratios with the fewest calculated values to ensure greater accuracy.

$$\frac{\sin 94^{\circ}}{15} \approx \frac{\sin 33^{\circ}}{b}$$
Law of Sines
$$b \approx \frac{15 \sin 33^{\circ}}{\sin 94^{\circ}} \text{ or about 8.2}$$
Solve for *b*.

Therefore, the remaining measures of $\triangle ABC$ are $B \approx 33^\circ$, $C \approx 53^\circ$, and $b \approx 8.2$.

b. $a = 9, b = 11, A = 61^{\circ}$

Notice that *A* is acute and a < b because 9 < 11. Find *h*.

Definition of sine

 $h = 11 \sin 61^\circ$ or about 9.6 $h = b \sin A$

Because a < h, no triangle can be formed with sides a = 9, b = 11, and $A = 61^{\circ}$. Therefore, this problem has no solution.

GuidedPractice

 $\sin 61^\circ = \frac{h}{11}$

3A. $a = 12, b = 8, B = 61^{\circ}$

3B.
$$a = 13, c = 26, A = 30^{\circ}$$

S

StudyTip

Make a Reasonable Sketch When solving triangles, a

reasonably accurate sketch can

help you determine whether your

answer is feasible. In your sketch,

check to see that the longest side

is opposite the largest angle and

that the shortest side is opposite

the smallest angle.

 $C = \sin^{-1}\left(\frac{12\sin 94^{\circ}}{15}\right) \text{ or about } 53^{\circ}$

61

Example 4 The Ambiguous Case-Two Solutions

Find two triangles for which $A = 43^{\circ}$, a = 25, and b = 28. Round side lengths to the nearest tenth and angle measures to the nearest degree.

A is acute, and $h = 28 \sin 43^\circ$ or about 19.1. Notice that a < b because 25 < 28, and a > h because 25 > 19.1. Therefore, two different triangles are possible with the given angle and side measures. Angle *B* will be acute, while angle *B'* will be obtuse.



Make a reasonable sketch of each triangle and apply the Law of Sines to find each solution. Start with the case in which *B* is acute.

Law of Sines

Solve for sin B.

Use a calculator.

Definition of sin-1

Solution 1 $\angle B$ is acute.

Find B.

 $\frac{\sin B}{28} = \frac{\sin 43^\circ}{25}$

 $\sin B = \frac{28 \sin 43^\circ}{25}$

 $\sin B \approx 0.7638$

 $B \approx \sin^{-1} 0.7638$ or about 50°

Find C.

TechnologyTip

Using \sin^{-1} Notice that when calculating \sin^{-1} of a ratio, your

calculator will never return two

possible angle measures because \sin^{-1} is a *function*. Also, your calculator will never return an

obtuse angle measure for sin-1

has a range of $-\frac{\pi}{2}$ to $\frac{\pi}{2}$ or -90° to 90° .

because the inverse sine function

 $C\approx 180^\circ-(43^\circ+50^\circ)\approx 87^\circ$

Apply the Law of Sines to find c.

 $\frac{\sin 87^{\circ}}{c} \approx \frac{\sin 43^{\circ}}{25}$ Law of Sines $c \approx \frac{25 \sin 87^{\circ}}{\sin 43^{\circ}} \text{ or about } 36.6$ Solve for *c*.

Solution 2 $\angle B'$ is obtuse.

Note that $m \angle CB'B \cong m \angle CBB'$. To find B', you need to find an obtuse angle with a sine that is also 0.7638. To do this, subtract the measure given by your calculator to the nearest degree, 50°, from 180°. Therefore, B' is approximately $180^\circ - 50^\circ$ or 130° .

Find C.

 $C \approx 180^{\circ} - (43^{\circ} + 130^{\circ}) \text{ or } 7^{\circ}$

Apply the Law of Sines to find *c*.

$$\frac{\sin 7^{\circ}}{c} \approx \frac{\sin 43^{\circ}}{25}$$
Law of Sines
$$c \approx \frac{25 \sin 7^{\circ}}{\sin 43^{\circ}} \text{ or about } 4.5$$
Solve for *c*.

Therefore, the missing measures for acute $\triangle ABC$ are $B \approx 50^\circ$, $C \approx 87^\circ$, and $c \approx 36.6$, while the missing measures for obtuse $\triangle AB'C$ are $B' \approx 130^\circ$, $C \approx 7^\circ$, and $c \approx 4.5$.

GuidedPractice

Find two triangles with the given angle measure and side lengths. Round side lengths to the nearest tenth and angle measures to the nearest degree.

4A. $A = 38^{\circ}, a = 8, b = 10$

4B.
$$A = 65^{\circ}, a = 55, b = 57$$



25

R

StudyTip

Law of Cosines Notice that the angle referenced in each equation of the Law of Cosines corresponds to the side length on the other side of the equation.

 $a^{2} = b^{2} + c^{2} - 2bc \cos A$ $b^{2} = a^{2} + c^{2} - 2ac \cos B$ $c^{2} = a^{2} + b^{2} - 2ab \cos C$ You can use the **Law of Cosines** to solve an oblique triangle for the remaining two cases: when you are given the measures of three sides (SSS) or the measures of two sides and their included angle (SAS).

KeyConcept Law of Cosines

In \triangle ABC, if sides with lengths *a*, *b*, and *c* are opposite angles with measures *A*, *B*, and *C*, respectively, then the following are true.

 $a² = b² + c² - 2bc \cos A$ $b² = a² + c² - 2ac \cos B$ $c² = a² + b² - 2ab \cos C$



You will derive the first formula for the Law of Cosines in Exercise 70.

Real-World Example 5 Apply the Law of Cosines (SSS)

HOCKEY When a hockey player attempts a shot, he is 20 feet from the left post of the goal and 24 feet from the right post, as shown. If a regulation hockey goal is 6 feet wide, what is the player's shot angle to the nearest degree?



StudyTip

Check for Reasonableness

Because a triangle can have at most one obtuse angle, it is wise to find the measure of the largest angle in a triangle first, which will be the angle opposite the longest side. If the largest angle is obtuse, then you know that the other two angles must be acute. If the largest angle is acute, the remaining two angles must still be acute. Since three side lenghts are given, you can use the Law of Cosines to find the player's shot angle, *A*.

 $a^2 = b^2 + c^2 - 2bc \cos A$ Law of Cosines $6^2 = 24^2 + 20^2 - 2(24)(20) \cos A$ a = 6, b = 24, and c = 20 $36 = 576 + 400 - 960 \cos A$ Simplify. $36 = 976 - 960 \cos A$ Add. $-940 = -960 \cos A$ Subtract 976 from each side. $\frac{940}{960} = \cos A$ Divide each side by -960. $\cos^{-1}\left(\frac{940}{960}\right) = A$ Use the \cos^{-1} function. $11.7^{\circ} \approx A$ Use a calculator.

So, the player's shot angle is about 12°.

GuidedPractice

5. HIKING A group of friends who are on a camping trip decide to go on a hike. According to the map shown, what is the angle that is formed by the two trails that lead to the camp?



Example 6 Apply the Law of Cosines (SAS) Solve $\triangle ABC$. Round side lengths to the nearest tenth and angle measures to the nearest degree. 65° Step 1 Use the Law of Cosines to find the missing side measure. $c^2 = a^2 + b^2 - 2ab \cos C$ Law of Cosines R $c^2 = 5^2 + 8^2 - 2(5)(8) \cos 65^\circ$ $a = 5, b = 8, and C = 65^{\circ}$ $c^2 \approx 55.19$ Use a calculator. $c \approx 7.4$ Take the positive square root of each side. Step 2 Use the Law of Sines to find a missing angle measure. $\frac{\sin A}{5} = \frac{\sin 65^{\circ}}{7.4}$ $\frac{\sin A}{2} = \frac{\sin C}{C}$ $\sin A = \frac{5\sin 65^\circ}{7.4}$ Multiply each side by 5. $A \approx 38^{\circ}$ Definition of sin⁻¹ **Step 3** Find the measure of the remaining angle. $B \approx 180^{\circ} - (65^{\circ} + 38^{\circ}) \text{ or } 77^{\circ}$ Therefore, $c \approx 7.4$, $A \approx 38^\circ$, and $B \approx 77^\circ$. **Guided**Practice **6.** Solve $\triangle HJK$ if $H = 34^\circ$, j = 7, and k = 10.

2 Find Areas of Oblique Triangles When the measures of all three sides of a triangle are known, the Law of Cosines can be used to prove Heron's Formula for the area of the triangle.





StudyTip

Semiperimeter The measure *s* used in Heron's Formula is called the *semiperimeter* of the triangle.

In the ambiguous case of the Law of Sines, you compared the length of *a* to the value $h = b \sin A$. In the triangle shown, *h* represents the length of the altitude to side *c* in $\triangle ABC$. You can use this expression for the height of the triangle to develop a formula for the area of the triangle.



Area =
$$\frac{1}{2}ch$$
 Formula for area of a triangle
= $\frac{1}{2}c(b \sin A)$ Replace *h* with *b* sin *A*.
= $\frac{1}{2}bc \sin A$ Simplify.

By a similar argument, you can develop the formulas

Area
$$=\frac{1}{2}ab\sin C$$
 and Area $=\frac{1}{2}ac\sin B$

Notice that in each of these formulas, the information needed to find the area of the triangle is the measures of two sides and the included angle.



Because the area of a triangle is constant, the formulas above can be written as one formula.

Area
$$=\frac{1}{2}bc\sin A = \frac{1}{2}ab\sin C = \frac{1}{2}ac\sin B$$

If the included angle measures 90°, notice that each formula simplifies to the formula for the area of a right triangle, $\frac{1}{2}$ (base)(height), because sin 90° = 1.



StudyTip Area of an Obtuse Triangle

This formula works for any type of triangle, including obtuse triangles. You will prove this in Lesson 5-3. Solve each triangle. Round to the nearest tenth, if necessary. (Examples 1 and 2)



- **7. GOLF** A golfer misses a 12-foot putt by putting 3° off course. The hole now lies at a 129° angle between the ball and its spot before the putt. What distance does the golfer need to putt in order to make the shot? (Examples 1 and 2)
- **8. ARCHITECTURE** An architect's client wants to build a home based on the architect Jon Lautner's Sheats-Goldstein House. The length of the patio will be 60 feet. The left side of the roof will be at a 49° angle of elevation, and the right side will be at an 18° angle of elevation. Determine the lengths of the left and right sides of the roof and the angle at which they will meet. (Examples 1 and 2)



- **9 TRAVEL** For the initial 90 miles of a flight, the pilot heads 8° off course in order to avoid a storm. The pilot then changes direction to head toward the destination for the remainder of the flight, making a 157° angle to the first flight course. (Examples 1 and 2)
 - **a.** Determine the total distance of the flight.
 - **b.** Determine the distance of a direct flight to the destination.

Find all solutions for the given triangle, if possible. If no solution exists, write *no solution*. Round side lengths to the nearest tenth and angle measures to the nearest degree. (Example 3)

10. $a = 9, b = 7, A = 108^{\circ}$	11. $a = 14, b = 15, A = 117^{\circ}$
12. $a = 18, b = 12, A = 27^{\circ}$	13. $a = 35, b = 24, A = 92^{\circ}$
14. $a = 14, b = 6, A = 145^{\circ}$	15. $a = 19, b = 38, A = 30^{\circ}$
16. $a = 5, b = 6, A = 63^{\circ}$	17. $a = 10, b = \sqrt{200}, A = 45^{\circ}$

18. SKIING A ski lift rises at a 28° angle during the first 20 feet up a mountain to achieve a height of 25 feet, which is the height maintained during the remainder of the ride up the mountain. Determine the length of cable needed for this initial rise. (Example 3)



Find two triangles with the given angle measure and side lengths. Round side lengths to the nearest tenth and angle measures to the nearest degree. (Example 4)

19. $A = 39^{\circ}, a = 12, b = 17$	20. $A = 26^{\circ}, a = 5, b = 9$
21. $A = 61^\circ, a = 14, b = 15$	22. <i>A</i> = 47°, <i>a</i> = 25, <i>b</i> = 34
23. $A = 54^\circ, a = 31, b = 36$	24. $A = 18^{\circ}, a = 8, b = 13$

25. BROADCASTING A radio tower located 38 miles along Industrial Parkway transmits radio broadcasts over a 30-mile radius. Industrial Parkway intersects the interstate at a 41° angle. How far along the interstate can vehicles pick up the broadcasting signal? (Example 4)



26. BOATING The light from a lighthouse can be seen from an 18-mile radius. A boat is anchored so that it can just see the light from the lighthouse. A second boat is located 25 miles from the lighthouse and is headed straight toward it, making a 44° angle with the lighthouse and the first boat. Find the distance between the two boats when the second boat enters the radius of the lighthouse light. (Example 4)

Solve each triangle. Round side lengths to the nearest tenth and angle measures to the nearest degree. (Examples 5 and 6)

- **27.** $\triangle ABC$, if $A = 42^\circ$, b = 12, and c = 19
- **28.** $\triangle XYZ$, if x = 5, y = 18, and z = 14
- **29.** $\triangle PQR$, if $P = 73^{\circ}$, q = 7, and r = 15
- **30.** $\triangle JKL$, if $J = 125^{\circ}$, k = 24, and $\ell = 33$
- **31.** $\triangle RST$, if r = 35, s = 22, and t = 25
- **32.** \triangle *FGH*, if *f* = 39, *g* = 50, and *h* = 64
- **33.** $\triangle BCD$, if $B = 16^{\circ}$, c = 27, and d = 3
- **34.** $\triangle LMN$, if $\ell = 12$, m = 4, and n = 9

- **35. AIRPLANES** During her shift, a pilot flies from Columbus to Atlanta, a distance of 448 miles, and then on to Phoenix, a distance of 1583 miles. From Phoenix, she returns home to Columbus, a distance of 1667 miles. Determine the angles of the triangle created by her flight path. (Examples 5 and 6)
- **36. CATCH** Lola rolls a ball on the ground at an angle of 23° to the right of her dog Buttons. If the ball rolls a total distance of 48 feet, and she is standing 30 feet away, how far will Buttons have to run to retrieve the ball? (Examples 5 and 6)

Use Heron's Formula to find the area of each triangle. Round to the nearest tenth. (Example 7)

- **37.** x = 9 cm, y = 11 cm, z = 16 cm
- **38.** x = 29 in., y = 25 in., z = 27 in.
- **39.** x = 58 ft, y = 40 ft, z = 63 ft
- **40.** x = 37 mm, y = 10 mm, z = 34 mm
- **41.** x = 8 yd, y = 15 yd, z = 8 yd
- **42.** *x* = 133 mi, *y* = 82 mi, *z* = 77 mi
- **43.** LANDSCAPING The Steele family wants to expand their backyard by purchasing a vacant lot adjacent to their property. To get a rough measurement of the area of the lot, Mr. Steele counted the steps needed to walk around the border and diagonal of the lot. (Example 7)



- **a.** Estimate the area of the lot in steps.
- **b.** If Mr. Steele measured his step to be 1.8 feet, determine the area of the lot in square feet.
- **44. DANCE** During a performance, a dancer remained within a triangular area of the stage. (Example 7)



- **a.** Find the area of stage used in the performance.
- **b.** If the stage is 250 square feet, determine the percentage of the stage used in the performance.

Find the area of each triangle to the nearest tenth. (Example 8)

- **45.** $\triangle ABC$, if $A = 98^\circ$, b = 13 mm, and c = 8 mm
- **46.** $\triangle JKL$, if $L = 67^{\circ}$, j = 11 yd, and k = 24 yd
- **47.** $\triangle RST$, if $R = 35^{\circ}$, s = 42 ft, and t = 26 ft
- **48.** $\triangle XYZ$, if $Y = 124^{\circ}$, x = 16 m, and z = 18 m
- **49.** $\triangle FGH$, if $F = 41^{\circ}$, g = 22 in., and h = 36 in.
- **50.** $\triangle PQR$, if $Q = 153^{\circ}$, p = 27 cm, and r = 21 cm

51. DESIGN A free-standing art project requires a triangular support piece for stability. Two sides of the triangle must measure 18 and 15 feet in length and a nonincluded angle must measure 42°. If support purposes require the triangle to have an area of at least 75 square feet, what is the measure of the third side? (Example 8)

Use Heron's Formula to find the area of each figure. Round answers to the nearest tenth.



56. ZIP LINES A tourist attraction currently has its base connected to a tree platform 150 meters away by a zip line. The owners now want to connect the base to a second platform located across a canyon and then connect the platforms to each other. The bearings from the base to each platform and from platform 1 to platform 2 are given. Find the distances from the base to platform 2 and from platform 1 to platform 2.



57. LIGHTHOUSES The bearing from the South Bay lighthouse to the Steep Rock lighthouse 25 miles away is N 28° E. A small boat in distress spotted off the coast by each lighthouse has a bearing of N 50° W from South Bay and S 80° W from Steep Rock. How far is each tower from the boat?



connectED.mcgraw-hill.com

290

Find the area of each figure. Round answers to the nearest tenth.



62. BRIDGE DESIGN In the figure below, $\angle FDE = 45^{\circ}$, $\angle CED = 55^{\circ}$, $\angle FDE \cong \angle FGE$, *B* is the midpoint of *AC*, and $DE \cong EG$. If AD = 4 feet, DE = 12 feet, and CE = 14 feet, find *BF*.



63. BUILDINGS Barbara wants to know the distance between the tops of two buildings *R* and *S*. On the top of her building, she measures the distance between the points *T* and *U* and finds the given angle measures. Find the distance between the two buildings.



64. DRIVING After a high school football game, Della left the parking lot traveling at 35 miles per hour in the direction N 55° E. If Devon left 20 minutes after Della at 45 miles per hour in the direction S 10° W, how far apart are Devon and Della an hour and a half after Della left?



H.O.T. Problems Use Higher-Order Thinking Skills

- **65. ERROR ANALYSIS** Monique and Rogelio are solving an acute triangle in which $\angle A = 34^\circ$, a = 16, and b = 21. Monique thinks that the triangle has one solution, while Rogelio thinks that the triangle has no solution. Is either of them correct? Explain your reasoning.
- **66.** WRITING IN MATH Explain the different circumstances in which you would use the Law of Cosines, the Law of Sines, the Pythagorean Theorem, and the trigonometric ratios to solve a triangle.
- **67. REASONING** Why does an obtuse measurement appear on the graphing calculator for inverse cosine while negative measures appear for inverse sine?
- **68. PROOF** Show for a given rhombus with a side length of *s* and an included angle of θ that the area can be found with the formula $A = s^2 \sin \theta$.
- 69. PROOF Derive the Law of Sines.
- 70. **PROOF** Consider the figure below.



- **a.** Use the figure and hints below to derive the first formula $a^2 = b^2 + c^2 2bc \cos A$ in the Law of Cosines.
 - Use the Pythagorean Theorem for $\triangle DBC$.
 - In $\triangle ADB$, $c^2 = x^2 + h^2$.
 - $\cos A = \frac{x}{c}$
- **b.** Explain how you would go about deriving the other two formulas in the Law of Cosines.
- **CHALLENGE** A satellite is orbiting 850 miles above Mars and is now positioned directly above one of the poles. The radius of Mars is 2110 miles. If the satellite was positioned at point *X* 14 minutes ago, approximately how many hours does it take for the satellite to complete a full orbit, assuming that it travels at a constant rate around a circular orbit?



72. WRITING IN MATH Describe why solving a triangle in which h < a < b using the Law of Sines results in two solutions. Is this also true when using the Law of Cosines? Explain your reasoning.

Spiral Review

Find the exact value of each expression, if it exists. (Lesson 4-6)

73.
$$\cos^{-1} - \frac{1}{2}$$
 74. $\sin^{-1} \frac{\sqrt{2}}{2}$ **75.** $\arctan 1$

76. $\sin^{-1}\frac{\sqrt{3}}{2}$

Identify the damping factor f(x) of each function. Then use a graphing calculator to sketch the graphs of f(x), -f(x), and the given function in the same viewing window. Describe the behavior of the graph. (Lesson 4-5)

77.
$$y = -2x \sin x$$
 78. $y = \frac{3}{5}x \cos x$ **79.** $y = (x - 1)^2 \sin x$ **80.** $y = -4x^2 \cos x$

- 81. CARTOGRAPHY The distance around Earth along a given latitude can be found using $C = 2\pi r \cos L$, where *r* is the radius of Earth and *L* is the latitude. The radius of Earth is approximately 3960 miles. Make a table of values for the latitude and corresponding distance around Earth that includes $L = 0^{\circ}$, 30° , 45° , 60° , and 90° . Use the table to describe the distances along the latitudes as you go from 0° at the equator to 90° at a pole. (Lesson 4-3)
- 82. RADIOACTIVITY A scientist starts with a 1-gram sample of lead-211. The amount of the sample remaining after various times is shown in the table below. (Lesson 3-5)

Time (min)	10	20	30	40
Pb-211 present (g)	0.83	0.68	0.56	0.46

- **a.** Find an exponential regression equation for the amount *y* of lead as a function of time *x*.
- **b.** Write the regression equation in terms of base *e*.
- **c.** Use the equation from part **b** to estimate when there will be 0.01 gram of lead-211 present.

Write a polynomial function of least degree with real coefficients in standard form that has the given zeros. (Lesson 2-4)

83. $-1, 1, 5$ 84. $-2, -0.5, 4$ 85. $-3, -2i, 2i$	86. $-5i, -i, i, 5i$
---	-----------------------------



89. FREE RESPONSE The pendulum at the right moves according to $\theta = \frac{1}{4} \cos 12t$, where θ is the angular displacement in radians and *t* is the time in



- a. Set the mode to radians and graph the function
- **b.** What are the period, amplitude, and frequency of the function? What do they mean in the context of
- c. What is the maximum angular displacement of the pendulum in degrees?
- d. What does the midline of the graph represent?
- e. At what times is the pendulum displaced 5 degrees?

Chapter Summary

KevConcepts

Right Triangle Tr	igonometry (Lesson 4-1)	
$\sin \theta = \frac{\text{opp}}{1}$	$\cos \theta = \frac{\mathrm{adj}}{\mathrm{adj}}$	$\tan \theta = \frac{\operatorname{opp}}{\operatorname{opp}}$
hyp	hyp	adj
$\csc \theta = \frac{hyp}{hyp}$	sec $\theta = \frac{hyp}{hyp}$	$\cot \theta - \frac{\operatorname{adj}}{\operatorname{adj}}$
$cord = \frac{1}{opp}$	$\frac{30000}{adj}$	$cor v = \frac{1}{opp}$

Degrees and Radians (Lesson 4-2)

- •
- To convert from radians to degrees, multiply by $\frac{180^{\circ}}{\pi}$ radians
- $v = \frac{s}{t}$, where s is the arc length traveled Linear speed: durina time t
- Angular speed: $\omega = \frac{\theta}{t}$, where θ is the angle of rotation (in radians) moved during time t

Trigonometric Functions on the Unit Circle (Lesson 4-3)

- For an angle θ in radians containing (x, y), $\cos \theta = \frac{x}{r}$, $\sin \theta = \frac{y}{r}$, and $\tan \theta = \frac{y}{x}$, where $r = \sqrt{x^2 + y^2}$.
- For an angle *t* containing (*x*, *y*) on the unit circle, $\cos \theta = x$, $\sin \theta = y$, and $\tan \theta = \frac{y}{y}$.

Graphing Sine and Cosine Functions (Lesson 4-4)

• A sinusoidal function is of the form $y = a \sin(bx + c) + d$ or $y = a \cos(bx + c) + d$, where amplitude = |a|, period $= \frac{2\pi}{|b|}$ frequency $= \frac{|b|}{2\pi}$, phase shift $= -\frac{c}{|b|}$, and vertical shift = d.

Graphing Other Trigonometric Functions (Lesson 4-5)

• A damped trigonometric function is of the form $y = f(x) \sin bx$ or $y = f(x) \cos bx$, where f(x) is the damping factor.

Inverse Trigonometric Functions (Lesson 4-6)

- $y = \sin^{-1} x$ iff $\sin y = x$, for $-1 \le x \le 1$ and $-\frac{\pi}{2} \le y \le \frac{\pi}{2}$.
- $y = \cos^{-1} x$ iff $\cos y = x$, for $-1 \le x \le 1$ and $0 \le y \le \pi$.
- $y = \tan^{-1} x$ iff $\tan y = x$, for $-\infty < x < \infty$ and $-\frac{\pi}{2} < y < \frac{\pi}{2}$.

The Law of Sines and the Law of Cosines (Lesson 4-7)

Let $\triangle ABC$ be any triangle.

- The Law of Sines:
- The Law of Sines: $\frac{\sin A}{a} = \frac{\sin B}{b} = \frac{\sin C}{c}$ The Law of Cosines: $a^2 = b^2 + c^2 2bc \cos A$

$$b^{2} = a^{2} + c^{2} - 2ac \cos B$$

 $c^{2} = a^{2} + b^{2} - 2ab \cos C$

KevVocabularv

amplitude (p. 257) angle of depression (p. 224) angle of elevation (p. 224) angular speed (p. 236) circular function (p. 248) cosecant (p. 220) cosine (p. 220) cotangent (p. 220) coterminal angles (p. 234) damped trigonometric function (p. 275) damped wave (p. 275) damping factor (p. 275) frequency (p. 260) initial side (p. 231) inverse trigonometric function (p. 223) Law of Cosines (p. 295) Law of Sines (p. 291) linear speed (p. 236) midline (p. 262)

oblique triangles (p. 291) period (p. 250) periodic function (p. 250) phase shift (p. 261) quadrantal angle (p. 243) radian (p. 232) reciprocal function (p. 220) reference angle (p. 244) secant (p. 220) sector (p. 237) sine (p. 220) sinusoid (p. 256) standard position (p. 231) tangent (p. 220) terminal side (p. 231) trigonometric functions (p. 220) trigonometric ratios (p. 220) unit circle (p. 247) vertical shift (p. 262)

VocabularyCheck

State whether each sentence is true or false. If false, replace the underlined term to make a true sentence.

- 1. The sine of an acute angle in a right triangle is the ratio of the lengths of its opposite leg to the hypotenuse.
- 2. The secant ratio is the reciprocal of the sine ratio.
- 3. An angle of elevation is the angle formed by a horizontal line and an observer's line of sight to an object below the line.
- 4. The radian measure of an angle is equal to the ratio of the length of its intercepted arc to the radius.
- 5. The rate at which an object moves along a circular path is called its linear speed.
- **6.** 0°, π , and $-\frac{\pi}{2}$ are examples of <u>reference angles</u>.
- 7. The period of the graph of $y = 4 \sin 3x$ is 4.
- **8.** For $f(x) = \cos bx$, as *b* increases, the frequency decreases.
- **9.** The range of the arcsine function is $[0, \pi]$.
- 10. The Law of Sines can be used to determine unknown side lengths or angle measures of some triangles.

Lesson-by-Lesson Review



Degrees and Radians (pp. 231-241)

Write each degree measure in radians as a multiple of π and each radian measure in degrees.

17.	135°	18.	450°
19.	$\frac{7\pi}{4}$	20.	$\frac{13\pi}{10}$

19.
$$\frac{7\pi}{4}$$
 20.

Identify all angles coterminal with the given angle. Then find and draw one positive and one negative angle coterminal with the given angle.

10 m

21.
$$342^{\circ}$$
 22. $-\frac{\pi}{6}$



Example 2

Identify all angles coterminal with $\frac{5\pi}{12}$. Then find and draw one positive and one negative coterminal angle.

All angles measuring $\frac{5\pi}{12} + 2n\pi$ are coterminal with a $\frac{5\pi}{12}$ angle. Let n = 1 and -1.

$$\frac{5\pi}{6} + 2\pi(1) = \frac{17\pi}{6}$$

$$\frac{5\pi}{6} - 2\pi(-1) = -\frac{7\pi}{6}$$







Trigonometric Functions on the Unit Circle (pp. 242–253)

Example 3 Sketch each angle. Then find its reference angle. **25.** 240° **26.** 75° Let $\cos \theta = \frac{5}{13}$, where $\sin \theta < 0$. Find the exact values of the **27.** $-\frac{3\pi}{4}$ **28.** $\frac{11\pi}{18}$ five remaining trigonometric functions of θ . Since $\cos \theta$ is positive and $\sin \theta$ is negative, θ lies in Quadrant IV. Find the exact values of the five remaining trigonometric This means that the *x*-coordinate of a point on the terminal side of θ is positive and the y-coordinate is negative. functions of θ . **29.** $\cos \theta = \frac{2}{\epsilon}$, where $\sin \theta > 0$ and $\tan \theta > 0$ Since $\cos \theta = \frac{x}{r} = \frac{5}{13}$, use x = 5 and r = 13 to find y. **30.** tan $\theta = -\frac{3}{4}$, where sin $\theta > 0$ and cos $\theta < 0$ $y = \sqrt{r^2 - x^2}$ Pythagorean Theorem $=\sqrt{169-25}$ or 12 r=13 and x=5**31.** sin $\theta = -\frac{5}{13}$, where $\cos \theta > 0$ and $\cot \theta < 0$ $\sin \theta = \frac{y}{r} \operatorname{or} \frac{12}{13}$ $\tan \theta = \frac{y}{x} \operatorname{or} \frac{12}{5}$ $\operatorname{sec} \theta = \frac{r}{x} \operatorname{or} \frac{13}{5}$ **32.** cot $\theta = \frac{2}{2}$, where sin $\theta < 0$ and tan $\theta > 0$ $\csc \theta = \frac{r}{v} \operatorname{or} \frac{13}{12}$ $\cot \theta = \frac{x}{v} \operatorname{or} \frac{5}{12}$ Find the exact value of each expression. If undefined, write undefined. **34.** $\cot \frac{11\pi}{6}$ 33. sin 180° **36.** $\cos\left(-\frac{19\pi}{6}\right)$ 35. sec 450°

Graphing Sine and Cosine Functions (pp. 256–266)

Describe how the graphs of f(x) and g(x) are related. Then find the amplitude and period of g(x), and sketch at least one period of both functions on the same coordinate axes.

37.	$f(x) = \sin x$	38.	$f(x) = \cos x$
	$g(x) = 5 \sin x$		$g(x) = \cos 2x$
39.	$f(x) = \sin x$ $g(x) = \frac{1}{2}\sin x$	40.	$f(x) = \cos x$ $g(x) = -\cos \frac{1}{2}x$
	Z		- 3

State the amplitude, period, frequency, phase shift, and vertical shift of each function. Then graph two periods of the function.

41.	$y = 2 \cos(x - \pi)$	42. $y = -\sin 2x + 1$
43.	$y = \frac{1}{2} \cos\left(x + \frac{\pi}{2}\right)$	44. $y = 3 \sin \left(x + \frac{2\pi}{3} \right)$

Example 4

Amplitude: |a|

State the amplitude, period, frequency, phase shift, and vertical shift of $y = 4 \sin \left(x - \frac{\pi}{2}\right) - 4$. Then graph two periods of the function.

In this function,
$$a = 4$$
, $b = 1$, $c = -\frac{\pi}{2}$, and $d = -4$.

= |4| or 4 Period:
$$\frac{2\pi}{|b|} = \frac{2\pi}{|1|}$$
 or 2π

Frequency:
$$\frac{|b|}{2\pi} = \frac{|1|}{2\pi}$$
 or $\frac{1}{2\pi}$

Vertical shift:
$$d$$
 or -4

Frequency: $\frac{|b|}{2\pi} = \frac{|1|}{2\pi}$ or $\frac{1}{2\pi}$ Phase shift: $-\frac{c}{|b|} = -\frac{-\frac{\pi}{2}}{|1|}$ or $\frac{\pi}{2}$

First, graph the midline y = -4. Then graph $y = 4 \sin x$ shifted $\frac{\pi}{2}$ units to the right and 4 units down.





4-6 Inverse Trigonometric Functions (pp. 280–290)				
Find the exact value of each expression, if it exists.		Example 6		
53. sin ⁻¹ (-1)	54. $\cos^{-1} \frac{\sqrt{3}}{2}$	Find the exact value of $-\sqrt{3}$.		
55. $\tan^{-1}\left(-\frac{\sqrt{3}}{3}\right)$	56. arcsin 0	Find a point on the unit circle in the interval $\left(-\frac{\pi}{2}, \frac{\pi}{2}\right)$ with a tangent		
57. arctan (-1)	58. arccos $\frac{\sqrt{2}}{2}$	of $-\sqrt{3}$. When $t = -\frac{\pi}{3}$, tan $t = -\sqrt{3}$. Therefore, $arctan -\sqrt{3} = -\frac{\pi}{3}$		
59. $\sin^{-1}\left[\sin\left(-\frac{\pi}{3}\right)\right]$	60. $\cos^{-1} [\cos (-3\pi)]$	3.		

The Law of Sines and the Law of Cosines (pp. 291–301)

Find all solutions for the given triangle, if possible. If no solution exists, write *no solution*. Round side lengths to the nearest tenth and angle measurements to the nearest degree.

61. $a = 11, b = 6, A = 22^{\circ}$ **62.** $a = 9, b = 10, A = 42^{\circ}$

63. $a = 20, b = 10, A = 78^{\circ}$ **64.** $a = 2, b = 9, A = 88^{\circ}$

Solve each triangle. Round side lengths to the nearest tenth and angle measures to the nearest degree.

65. a = 13, b = 12, c = 8 **66.** $a = 4, b = 5, C = 96^{\circ}$

Example 7

Solve the triangle if a = 3, b = 4, and $A = 71^{\circ}$.

In the figure, $h = 4 \sin 71^\circ$ or about 3.8

Because $a \le h$, there is no triangle that can be formed with sides a = 3, b = 4, and $A = 71^{\circ}$. Therefore, this problem has no solution.



Applications and Problem Solving

- **67. CONSTRUCTION** A construction company is installing a three-foothigh wheelchair ramp onto a landing outside of an office. The angle of the ramp must be 4°. (Lesson 4-1)
 - a. What is the length of the ramp?
 - **b.** What is the slope of the ramp?
- **68. NATURE** For a photography project, Maria is photographing deer from a tree stand. From her sight 30 feet above the ground, she spots two deer in a straight line, as shown below. How much farther away is the second deer than the first? (Lesson 4-1)



- 69. FIGURE SKATING An Olympic ice skater performs a routine in which she jumps in the air for 2.4 seconds while spinning 3 full revolutions. (Lesson 4-2)
 - a. Find the angular speed of the figure skater.
 - **b.** Express the angular speed of the figure skater in degrees per minute.
- **70. TIMEPIECES** The length of the minute hand of a pocket watch is 1.5 inches. What is the area swept by the minute hand in 40 minutes? (Lesson 4-2)



- WORLD'S FAIR The first Ferris wheel had a diameter of 250 feet and took 10 minutes to complete one full revolution. (Lesson 4-3)
 - a. How many degrees would the Ferris wheel rotate in 100 seconds?
 - **b.** How far has a person traveled if he or she has been on the Ferris wheel for 7 minutes?
 - c. How long would it take for a person to travel 200 feet?

- 72. AIR CONDITIONING An air-conditioning unit turns on and off to maintain the desired temperature. On one summer day, the air conditioner turns on at 8:30 A.M. when the temperature is 80° Fahrenheit and turns off at 8:55 A.M. when the temperature is 74°. (Lesson 4-4)
 - **a.** Find the amplitude and period if you were going to use a trigonometric function to model this change in temperature, assuming that the temperature cycle will continue.
 - **b.** Is it appropriate to model this situation with a trigonometric function? Explain your reasoning.
- 73. TIDES In Lewis Bay, the low tide is recorded as 2 feet at 4:30 A.M., and the high tide is recorded as 5.5 feet at 10:45 A.M. (Lesson 4-4)
 - a. Find the period for the trigonometric model.
 - **b.** At what time will the next high tide occur?
- **74. MUSIC** When plucked, a bass string is displaced 1.5 inches, and its damping factor is 1.9. It produces a note with a frequency of 90 cycles per second. Determine the amount of time it takes the string's motion to be dampened so that -0.1 < y < 0.1. (Lesson 4-5)
- **75. PAINTING** A painter is using a 15-foot ladder to paint the side of a house. If the angle the ladder makes with the ground is less than 65°, it will slide out from under him. What is the greatest distance that the bottom of the ladder can be from the side of the house and still be safe for the painter? (Lesson 4-6)



- **76.** NAVIGATION A boat is 20 nautical miles from a port at a bearing 30° north of east. The captain sees a second boat and reports to the port that his boat is 15 nautical miles from the second boat, which is located due east of the port. Can port personnel be sure of the second boat's position? Justify your answer. (Lesson 4-7)
- 77. GEOMETRY Consider quadrilateral *ABCD*. (Lesson 4-7)
 - a. Find C.
 - **b.** Find the area of *ABCD*.



Practice Test

Find the value of x. Round to the nearest tenth, if necessary.



Find the measure of angle θ . Round to the nearest degree, if necessary.



5. MULTIPLE CHOICE What is the linear speed of a point rotating at an angular speed of 36 radians per second at a distance of 12 inches from the center of the rotation?

A	420 in./s	C	439 in./s
В	432 in./s	D	444 in./s

Write each degree measure in radians as a multiple of π and each radian measure in degrees.

6. 200° **7.** $-\frac{8\pi}{3}$

8. Find the area of the sector of the circle shown.



Sketch each angle. Then find its reference angle.

9.
$$165^{\circ}$$
 10. $\frac{21\pi}{13}$

Find the exact value of each expression.

11. $\sec \frac{7\pi}{6}$ 12. $\cos \pi$	(-240°)
--	---------

13. MULTIPLE CHOICE An angle θ satisfies the following inequalities: $\csc \theta < 0$, $\cot \theta > 0$, and $\sec \theta < 0$. In which quadrant does θ lie?

F	I	н	III
G	II	J	IV

State the amplitude, period, frequency, phase shift, and vertical shift of each function. Then graph two periods of the function.

14.
$$y = 4\cos\frac{x}{2} - 5$$
 15. $y = -\sin\left(x + \frac{\pi}{2}\right)$

16. TIDES The table gives the approximate times that the high and low tides occurred in San Azalea Bay over a 2-day period.

Tide	High 1	Low 1 High 2		Low 2	
Day 1	2:35 а.м.	8:51 а.м.	3:04 р.м.	9:19 р.м.	
Day 2	3:30 а.м.	9:48 а.м.	3:55 р.м.	10:20 р.м.	

- **a.** The tides can be modeled with a trigonometric function. Approximately what is the period of this function?
- **b.** The difference in height between the high and low tides is 7 feet. What is the amplitude of this function?
- c. Write a function that models the tides where t is measured in hours. Assume the function has no phase shift or vertical shift.

Locate the vertical asymptotes, and sketch the graph of each function.

17.
$$y = \tan\left(x + \frac{\pi}{4}\right)$$
 18. $y = \frac{1}{2}\sec 2x$

Find all solutions for the given triangle, if possible. If no solution exists, write *no solution*. Round side lengths to the nearest tenth and angle measurements to the nearest degree.

19.
$$a = 8, b = 16, A = 22^{\circ}$$
20. $a = 9, b = 7, A = 84^{\circ}$ **21.** $a = 3, b = 5, c = 7$ **22.** $a = 8, b = 10, C = 46^{\circ}$

Find the exact value of each expression, if it exists.

23.
$$\cos^{-1}\left(-\frac{\sqrt{3}}{2}\right)$$
 24. $\sin^{-1}\left(-\frac{1}{2}\right)$

25. NAVIGATION A boat leaves a dock and travels 45° north of west averaging 30 knots for 2 hours. The boat then travels directly west averaging 40 knots for 3 hours.



- **a.** How many nautical miles is the boat from the dock after 5 hours?
- **b.** How many degrees south of east is the dock from the boat's present position?

connectED.mcgraw-hill.com

30

HAPTER.

Connect to AP Calculus Related Rates

Objective

 Model and solve related rates problems. If air is being pumped into a balloon at a given rate, can we find the rate at which the volume of the balloon is expanding? How does the rate a company spends money on advertising affect the rate of its sales? *Related rates* problems occur when the rate of change for one variable can be found by *relating* that to rates of change for other variables.

Suppose two cars leave a point at the same time. One car is traveling 40 miles per hour due north, while the second car is traveling 30 miles per hour due east. How far apart are the two cars after 1 hour? 2 hours? 3 hours? We can use the formula d = rt and the Pythagorean Theorem to solve for these values.

 $d = \sqrt{40^2 + 30^2}$

In this situation, we know the rates of change for each car. What if we want to know the rate at which the distance between the two cars is changing?

Activity 1 Rate of Change

Two cars leave a house at the same time. One car travels due north at 35 miles per hour, while the second car travels due east at 55 miles per hour. Approximate the rate at which the distance between the two cars is changing.

105

35



- **Step 2** Write equations for the distance traveled by each car after *t* hours.
- **Step 3** Find the distance traveled by each car after 1, 2, 3, and 4 hours.
- **Step 4** Use the Pythagorean Theorem to find the distance between the two cars at each point in time.
- **Step 5** Find the average rate of change of the distance between the two cars for $1 \le t \le 2, 2 \le t \le 3$, and $3 \le t \le 4$.

Analyze the Results

- 1. Make a scatter plot displaying the total distance between the two cars. Let time t be the independent variable and total distance *d* be the dependent variable. Draw a line through the points.
- **2.** What type of function does the graph seem to model? How is your conjecture supported by the values found in Step 5?
- **3.** What would happen to the average rate of change of the distance between the two cars if one of the cars slowed down? sped up? Explain your reasoning.

The rate that the distance between the two cars is changing is *related* to the rates of the two cars. In calculus, problems involving related rates can be solved using *implicit differentiation*. However, before we can use advanced techniques of differentiation, we need to understand how the rates involved relate to one another. Therefore, the first step to solving any related rates problem should always be to model the situation with a sketch or graph and to write equations using the relevant values and variables.

Activity 2 Model Related Rates

A rock tossed into a still body of water creates a circular ripple that grows at a rate of 5 centimeters per second. Find the area of the circle after 3 seconds if the radius of the circle is 5 centimeters at t = 1.

Step 1 Make a sketch of the situation.

Step 2 Write an equation for the radius *r* of the circle after *t* seconds.

Step 3 Find the radius for t = 3, and then find the area.

Analyze the Results

- 4. Find an equation for the area *A* of the circle in terms of *t*.
- **5.** Find the area of the circle for t = 1, 2, 3, 4, and 5 seconds.
- 6. Make a graph of the values. What type of function does the graph seem to model?

You can use the difference quotient to calculate the rate of change for the area of the circle at a certain point in time.

Activity 3 Approximate Related Rate

Approximate the rate of change for the area of the circle in Activity 2.

Step 1 Substitute the expression for the area of the circle into the difference quotient.

$$m = \frac{\pi [5(t+h)]^2 - \pi (5t)^2}{h}$$

= 3= 2

Step 2 Approximate the rate of change of the circle at 2 seconds. Let h = 0.1, 0.01, and 0.001.

Step 3 Repeat Steps 1 and 2 for t = 3 seconds and t = 4 seconds.

Analyze the Results

- 7. What do the rates of change appear to approach for each value of *t*?
- 8. What happens to the rate of change of the area of the circle as the radius increases? Explain.
- 9. How does this approach differ from the approach you used in Activity 1 to find the rate of change for the distance between the two cars? Explain why this was necessary.

Model and Apply

- **10.** A 13-foot ladder is leaning against a wall so that the base of the ladder is exactly 5 feet from the base of the wall. If the bottom of the ladder starts to slide away from the wall at a rate of 2 feet per second, how fast is the top of the ladder sliding down the wall?
 - **a.** Sketch a model of the situation. Let *d* be the distance from the top of the ladder to the ground and *m* be the rate at which the top of the ladder is sliding down the wall.
 - **b.** Write an expression for the distance from the base of the ladder to the wall after *t* seconds.
 - **c.** Find an equation for the distance *d* from the top of the ladder to the ground in terms of *t* by substituting the expression found in part *b* into the Pythagorean Theorem.
 - **d**. Use the Pythagorean Theorem to find the distance *d* from the top of the ladder to the ground for *t* = 0, 1, 2, 3, 3.5, and 3.75.
 - e. Make a graph of the values. What type of function does the graph seem to model?
 - f. Use the difference quotient to approximate the rate of change *m* for the distance from the top of the ladder to the ground at t = 2. Let h = 0.1, 0.01, and 0.001. As h approaches 0, what do the values for *m* appear to approach?

StudyTip **Difference Quotient** Recall that the difference quotient for calculating the slope of the line tangent to the graph of f(x) at the point (x, f(x)) is $m = \frac{f(x+h) - f(x)}{h(x)}$

h

connectED.mcgraw-hill.com 309

Trigonometric Identities and Equations

CHAPTER

Then	Now		Nhy? 🔺			
In Chapter 4, you learned to graph trigonometric functions and to solve right and oblique triangles.	 In Chapter 5, you will: Use and verify trigonometric identities. Solve trigonometric equations. Use sum and difference identities to evaluate trigonometric expressions and solve equations. Use double-angle, power-reducing, half-angle, and product-sum identities to evaluate trigonometric expressions and solve equations. 		 BUSINESS Musicians tune their instruments by listening for a <i>l</i> which is an interference between two sound waves with slightly different frequencies. The sum of the sound waves can be represented using a trigonometric equation. PREREAD Using what you know about trigonometric functions, make a prediction of what you will learn in Chapter 5. 			
connectED.mcgra	w-hill.com Your Digita	I Math Portal				
Animation V	ocabulary eGlossary	Personal Tutor	Graphing Calculator	Audio	Self-Check Practice	Work
<u>R</u>	B	PT		1		

r a *beat,* ghtly

orksheets

on/Corbis